

MATH 122 – THIRD UNIT TEST

Friday, November 21, 2008.

NAME: _____

Sona
Akopian

A

Brian
Seguin

B

C

Paul
McKenney

D

E

Oleksii
Mostovyi

F

G

Jason
Rute

H

I

Lisa
Espig

J

K

Instructions:

1. Do not separate the pages of the exam. If any pages do become separated, write your name on them and point them out to your instructor when you turn in the exam.
2. Please read the instructions for each individual question carefully. One of the skills being tested on this exam is your ability to interpret questions, so instructors will not answer questions about exam problems during the exam.
3. Show an appropriate amount of work for each exam question so that graders can see your final answer **and** how you obtained it.
4. You may use your calculator on all exam questions except where otherwise indicated. However, if you are asked to find an *exact* value of a quantity that involves an integral then you should not use calculator integration for this.
5. If you use graphs or tables to obtain an answer (especially if you create the graphs or tables on your calculator), be certain to provide an explanation and a sketch of the graph to show how you obtained your answer.
6. If you evaluate an improper integral, be sure to use appropriate algebraic and limit notation.
7. Please **TURN OFF** all cell phones and pagers, and **REMOVE** all headphones.

Problem	Total	Score
1	15	
2	20	
3	30	
4	20	
5	15	
Total	100	

1. 15 Points. SHOW ALL WORK. NO WORK = NO CREDIT.

Consider the infinite series:

$$\sum_{k=1}^{\infty} \frac{3}{k^2 + 3k}.$$

- (a) **(4 points)** Use the Technique of Partial Fractions to write $\frac{3}{k^2 + 3k}$ as the difference of two simpler fractions. Your final answer should contain no unspecified quantities beyond k .

- (b) **(3 points)** Calculate the partial sum indicated below. Include at least **8 decimal places** in your answer.

$$\sum_{k=1}^5 \frac{3}{k^2 + 3k} =$$

- (c) **(4 points)** Find a simple formula that gives the N^{th} partial sum of the infinite series that is valid for $N > 2$. The only variable in your formula should be N .

- (d) **(4 points)** Find the **exact** sum of the infinite series $\sum_{k=1}^{\infty} \frac{3}{k^2 + 3k}$. If you express your final answer as a decimal, include at least **8 decimal places**.

2. 20 Points. SHOW ALL WORK. NO PARTIAL CREDIT WITHOUT WORK.

The function $f(x)$ has a Taylor series that converges to $f(x)$ for all x in the interval of convergence. The n^{th} derivative of $f(x)$ evaluated at $a = 10$ is:

$$f^{(n)}(a) = f^{(n)}(10) = \frac{(-1)^n \cdot (n-1)!}{2^n} \quad \text{for all } n \geq 1.$$

The value of the function at the point $a = 10$ is: $f(a) = f(10) = -1$.

- (a) **(4 points)** Write down the Taylor series of the function $f(x)$ about $a = 10$. To express your answer, use Σ notation.

Σ

- (b) **(8 points)** Find the **radius** of convergence for the Taylor series of $f(x)$ about $a = 10$. Show your work.

Continued on the next page.

The function $f(x)$ has a Taylor series that converges to $f(x)$ for all x in the interval of convergence. The n^{th} derivative of $f(x)$ evaluated at $a = 10$ is:

$$f^{(n)}(a) = f^{(n)}(10) = \frac{(-1)^n \cdot (n-1)!}{2^n} \quad \text{for all } n \geq 1.$$

The value of the function at the point $a = 10$ is: $f(a) = f(10) = -1$.

- (c) **(8 points)** Find the **interval** of convergence for the Taylor series of $f(x)$ about $a = 10$. Show your work and clearly indicate your final answer. No work = no credit.

3. 30 Points. SHOW YOUR WORK. NO WORK = NO CREDIT.

Determine the convergence or divergence of each of the following series. If you do not justify your answer, you will get zero credit, even if you circle the correct final answer.

In each case, demonstrate that your answer is correct in a step-by-step fashion using *an appropriate convergence test*. Be sure to explicitly state which convergence test you have used and show that it can be used with the series you are working on. Be careful to show all of your work. As the final part of your answer in each part, CIRCLE either CONVERGES or DIVERGES.

(a) (10 points)
$$\sum_{n=1}^{\infty} \frac{e^n}{n!}$$

JUSTIFICATION:

FINAL ANSWER (CIRCLE ONE): CONVERGES DIVERGES

Demonstrate that your answer is correct in a step-by-step fashion using *an appropriate convergence test*. Be sure to explicitly state which convergence test you have used and show that it can be used with the series you are working on. Be careful to show all of your work. As the final part of your answer in each part, CIRCLE either CONVERGES or DIVERGES.

(b) (10 points) $\sum_{n=1}^{\infty} n \cdot e^{-n^2}$

JUSTIFICATION:

FINAL ANSWER (CIRCLE ONE):

CONVERGES

DIVERGES

Demonstrate that your answer is correct in a step-by-step fashion using *an appropriate convergence test*. Be sure to explicitly state which convergence test you have used and show that it can be used with the series you are working on. Be careful to show all of your work. As the final part of your answer in each part, CIRCLE either CONVERGES or DIVERGES.

(c) (10 points)
$$\sum_{n=1}^{\infty} \frac{\sqrt{n}}{n^2 + 1}$$

JUSTIFICATION:

FINAL ANSWER (CIRCLE ONE):

CONVERGES

DIVERGES

4. 20 Points. SHOW YOUR WORK.

Consider the alternating series:

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{(2n)!}.$$

- (a) **(10 points)** Does the series converge absolutely, converge conditionally or diverge? Clearly state your answer and use a convergence test to demonstrate that your answer is correct.

Be sure to explicitly state which convergence test you have used and show that it can be used with the series you are working on. Be careful to show all of your work and clearly state your final conclusion.

Continued on the next page.

The alternating series from the previous page is:

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{(2n)!}.$$

- (b) **(6 points)** Let S represent the sum of the series. Suppose that S , the sum of the series, is approximated by the N^{th} partial sum:


$$S_N = \sum_{n=1}^N \frac{(-1)^n}{(2n)!}.$$

What is the smallest value of N that could be used to approximate S by S_N and an error of less than 0.0001?

- (c) **(4 points)** Approximate the sum of the series $\sum_{n=1}^{\infty} \frac{(-1)^n}{(2n)!}$ with an error of less than 0.0001. Include at least **8 decimal places** in your answer.

5. 15 Points. 3 POINTS FOR EACH CORRECT RESPONSE.

The diagram shows the graphs of four functions, $f(x)$, $g(x)$, $h(x)$ and $k(x)$.



The formulas given below are Taylor polynomials of degree 2 centered at zero ($a = 0$). Match the Taylor polynomials with the functions. If you believe that a particular Taylor polynomial could not possibly correspond to any of the functions $f(x)$, $g(x)$, $h(x)$ or $k(x)$ write “**NONE**” in the space provided beside it.

- | | | |
|-----|-------------------------|-------------------------------|
| (a) | $P_2(x) = 1 + x + x^2$ | Corresponding function: _____ |
| (b) | $P_2(x) = -1 + x + x^2$ | Corresponding function: _____ |
| (c) | $P_2(x) = 1 - x + x^2$ | Corresponding function: _____ |
| (d) | $P_2(x) = -1 + x - x^2$ | Corresponding function: _____ |
| (e) | $P_2(x) = -1 - x + x^2$ | Corresponding function: _____ |