


**Department of Mathematics**  
**Carnegie Mellon University**

21-301 Combinatorics, Fall 2010: Test 4

Name: \_\_\_\_\_

Problem	Points	Score
1	40	
2	40	
3	20	
Total	100	

**Q1: (40pts)**



How many ways are there of  $k$ -coloring the squares of the above picture if the group acting is  $e_0, e_2, p, q$  where  $e_j$  is rotation by  $2\pi j/4$  and  $p, q$  are horizontal and vertical reflections.

(All small squares are meant to be of the same size here).

**Solution**

$$|Fix(g)| \begin{array}{ccccc} g & e_0 & e_2 & p & q \\ k^{17} & k^9 & k^{12} & k^{12} & k^{12} \end{array}$$

So the total number of colorings is

$$\frac{k^{17} + k^9 + k^{12} + k^{12}}{4}.$$

**Q2: (40pts)**

Consider the following take-away game: There is a pile of  $n$  chips. A move consists of removing 1 or 4 chips. Determine the Sprague-Grundy numbers  $g(n)$  for  $n \geq 0$  and prove that they are what you claim.

**Solution:** After looking at the first few numbers 0, 1, 0, 1, 2, 0, 1, 0, 1, 2, ... one sees that

$$g(n) = \begin{cases} 0 & n = 0, 2 \pmod{5} \\ 1 & n = 1, 3 \pmod{5} \\ 2 & n = 4 \pmod{5} \end{cases}$$

We verify this by induction. It is true for  $n \leq 10$  by inspection. For  $n > 10$  we have that if  $n = 5m + s$  then

$$g(n) = \text{mex}\{g(n-1), g(n-4)\} = \text{mex}\{g(5(m-1)+s+4), g(5(m-1)+s+1)\}$$

So, by induction

$$g(n) = \begin{cases} \text{mex}\{g(5(m-1)+4), g(5(m-1)+1)\} = \text{mex}\{2, 1\} = 0 & s = 0 \\ \text{mex}\{g(5m), g(5(m-1)+2)\} = \text{mex}\{0, 0\} = 1 & s = 1 \\ \text{mex}\{g(5m+1), g(5(m-1)+3)\} = \text{mex}\{1, 1\} = 0 & s = 2 \\ \text{mex}\{g(5m+2), g(5(m-1)+4)\} = \text{mex}\{0, 2\} = 1 & s = 3 \\ \text{mex}\{g(5m+3), g(5m)\} = \text{mex}\{0, 1\} = 2 & s = 4 \end{cases}$$

The result follows by induction.

**Q3: (20pts)**

In the game Split Nim a player removes chips from a non-empty pile and then if desired, has the further option of splitting the reduced pile into two non-empty piles (if the reduced pile has more than one chip). Show that Split Nim has the same N and P positions as ordinary Nim.

**Solution:** We prove this by induction on the total number  $t$  of chips.  $t = 0$  is a P position in both games.

Now suppose that  $t > 0$  and the position is an N position for Nim. If the player uses regular Nim strategy then the resulting position is a P position for Nim and by induction this is a P position for Split Nim.

Suppose then that  $t > 0$  and the position is a P position for Nim. Suppose that the pile sizes are  $p_1, p_2, \dots, p_k$ . Suppose that the player first removes chips to leave  $p'_1$  chips in the first pile. We know that  $p'_1 \oplus p_2 \oplus \dots \oplus p_k \neq 0$  is an N position for Split Nim by induction.

So, suppose that the player now splits the first pile into two piles of size  $a, b$ . We will argue that  $a \oplus b \oplus p_2 \oplus \dots \oplus p_k \neq 0$ . This is an N position for Nim and it will be an N position for Split Nim by induction. Suppose to the contrary that  $a \oplus b \oplus p_2 \oplus \dots \oplus p_k = 0$ . We will argue that  $c = a \oplus b \leq a + b$ . It follows that the previous position was in fact an N position for Nim, since the player could have removed  $p_1 - c$  chips and left a P position.

But if  $a = \sum_i a_i 2^i$  and  $b = \sum_i b_i 2^i$  then

$$a + b - (a \oplus b) = \sum_i (a_i + b_i - (a_i \oplus b_i)) 2^i \geq 0.$$