




## Trees



A *tree* is a graph which is

**(a)** Connected and

**(b)** has no cycles (*acyclic*).


**Lemma 1** *Let the components of  $G$  be*

*$C_1, C_2, \dots, C_r$ , Suppose  $e = (u, v) \notin E, u \in C_i, v \in C_j$ .*


**(a)**  $i = j \Rightarrow \omega(G + e) = \omega(G)$ .

**(b)**  $i \neq j \Rightarrow \omega(G + e) = \omega(G) - 1$ .

(a)



(b)



**Proof** Every path  $P$  in  $G + e$  which is not in  $G$  must contain  $e$ . Also,

$$\omega(G + e) \leq \omega(G).$$

Suppose

$$(x = u_0, u_1, \dots, u_k = u, u_{k+1} = v, \dots, u_\ell = y)$$

is a path in  $G + e$  that uses  $e$ . Then clearly  $x \in C_i$  and  $y \in C_j$ .

(a) follows as now no new relations  $x \sim y$  are added.

(b) Only possible new relations  $x \sim y$  are for  $x \in C_i$  and  $y \in C_j$ . But  $u \sim v$  in  $G + e$  and so  $C_i \cup C_j$  becomes (only) new component.  $\square$

**Lemma 2**  $G = (V, E)$  is acyclic (forest) with (tree) components  $C_1, C_2, \dots, C_k$ .  $|V| = n$ .  $e = (u, v) \notin E$ ,  $u \in C_i$ ,  $v \in C_j$ .


**(a)**  $i = j \Rightarrow G + e$  contains a cycle.

**(b)**  $i \neq j \Rightarrow G + e$  is acyclic and has one less component.


**(c)**  $G$  has  $n - k$  edges.

(a)  $u, v \in C_i$  implies there exists a path  
 $(u = u_0, u_1, \dots, u_\ell = v)$  in  $G$ .

So  $G + e$  contains the cycle  $u_0, u_1, \dots, u_\ell, u_0$ .




(a)



Suppose  $G + e$  contains the cycle  $C$ .  $e \in C$  else  $C$  is a cycle of  $G$ .

$$C = (u = u_0, u_1, \dots, u_\ell = v, u_0).$$

But then  $G$  contains the path  $(u_0, u_1, \dots, u_\ell)$  from  $u$  to  $v$  – contradiction.



The drop in the number of components follows from Lemma 1.

The rest of the lemma follows from

(c) Suppose  $E = \{e_1, e_2, \dots, e_r\}$  and  $G_i = (V, \{e_1, e_2, \dots, e_i\})$  for  $0 \leq i \leq r$ .

**Claim:**  $G_i$  has  $n - i$  components.

Induction on  $i$ .

$i = 0$ :  $G_0$  has no edges.

$i > 0$ :  $G_{i-1}$  is acyclic and so is  $G_i$ . It follows from part (a) that  $e_i$  joins vertices in distinct components of  $G_{i-1}$ . It follows from (b) that  $G_i$  has one less component than  $G_{i-1}$ .

**End of proof of claim**

Thus  $r = n - k$  (we assumed  $G$  had  $k$  components).

□

**Corollary 1** *If a tree  $T$  has  $n$  vertices then*

**(a)** *It has  $n - 1$  edges.*

**(b)** *It has at least 2 vertices of degree 1, ( $n \geq 2$ ).*

**Proof** (a) is part (c) of previous lemma.  $k = 1$  since  $T$  is connected.

(b) Let  $s$  be the number of vertices of degree 1 in  $T$ . There are no vertices of degree 0 – these would form separate components. Thus

$$2n - 2 = \sum_{v \in V} d_T(v) \geq 2(n - s) + s.$$

So  $s \geq 2$ . □



**Theorem 1** *Suppose  $|V| = n$  and  $|E| = n - 1$ . The following three statements become equivalent.*

**(a)**  *$G$  is connected.*

**(b)**  *$G$  is acyclic.*

**(c)**  *$G$  is a tree.*


**Proof** Let  $E = \{e_1, e_2, \dots, e_{n-1}\}$  and  $G_i = (V, \{e_1, e_2, \dots, e_i\})$  for  $0 \leq i \leq n - 1$ .

(a)  $\Rightarrow$  (b):  $G_0$  has  $n$  components and  $G_{n-1}$  has 1 component. Addition of each edge  $e_i$  must reduce the number of components by 1 – Lemma 1(b). Thus  $G_{i-1}$  acyclic implies  $G_i$  is acyclic – Lemma 2(b). (b) follows as  $G_0$  is acyclic.

(b)  $\Rightarrow$  (c): We need to show that  $G$  is connected. Since  $G_{n-1}$  is acyclic,  $\omega(G_i) = \omega(G_{i-1}) - 1$  for each  $i$  – Lemma 2(b). Thus  $\omega(G_{n-1}) = 1$ .


(c)  $\Rightarrow$  (a): trivial.

**Corollary 2** *If  $v$  is a vertex of degree 1 in a tree  $T$  then  $T - v$  is also a tree.*



**Proof** Suppose  $T$  has  $n$  vertices and  $n$  edges. Then  $T - v$  has  $n - 1$  vertices and  $n - 2$  edges. It is acyclic and so must be a tree.  $\square$

## Cut edges




$e$  is a *cut edge* of  $G$  if  $\omega(G - e) > \omega(G)$ .

**Theorem 2**  $e = (u, v)$  is a *cut edge* iff  $e$  is not on any *cycle* of  $G$ .

**Proof**  $\omega$  increases iff there exist  $x \sim y \in V$  such that all walks from  $x$  to  $y$  use  $e$ .

Suppose there is a cycle  $(u, P, v, u)$  containing  $e$ . Then if  $W = x, W_1, u, v, W_2, y$  is a walk from  $x$  to  $y$  using  $e$ ,  $x, W_1, P, W_2, y$  is a walk from  $x$  to  $y$  that doesn't use  $e$ . Thus  $e$  is not a cut edge.



If  $e$  is not a cut edge then  $G - e$  contains a path  $P$  from  $u$  to  $v$  ( $u \sim v$  in  $G$  and relations are maintained after deletion of  $e$ ). So  $(v, u, P, v)$  is a cycle containing  $e$ .

□

**Corollary 3** *A connected graph is a tree iff every edge is a cut edge.*

**Corollary 4** *Every finite connected graph  $G$  contains a spanning tree.*

**Proof** Consider the following process: starting with  $G$ ,

1. If there are no cycles – **stop**.
2. If there is a cycle, delete an edge of a cycle.

Observe that (i) the graph remains connected – we delete edges of cycles. (ii) the process must terminate as the number of edges is assumed finite.

On termination there are no cycles and so we have a connected acyclic spanning subgraph i.e. we have a spanning tree.  $\square$

## Alternative Construction

Let  $E = \{e_1, e_2, \dots, e_m\}$ .

begin

$T := \emptyset$

for  $i = 1, 2, \dots, m$  do

begin

if  $T + e_i$  does not contain a cycle

then  $T \leftarrow T + e_i$

end

Output  $T$

end

**Lemma 3** *If  $G$  is connected then  $(V, T)$  is a spanning tree of  $G$ .*


**Proof** Clearly  $T$  is acyclic. Suppose it is not connected and has components  $C_1, C_2, \dots, C_k$ ,  $k \geq 2$ . Let  $D = C_2 \cup \dots \cup C_k$ . Then  $G$  has no edges joining  $C_1$  and  $D$  – contradiction. (The first  $C_1 : D$  edge found by the algorithm would have been added.)



**Theorem 3** *Let  $T$  be a spanning tree of  $G = (V, E)$ ,  $|V| = n$ . Suppose  $e = (u, v) \in E \setminus T$ .*

**(a)**  *$T + e$  contains a unique cycle  $C(T, e)$ .*

**(b)**  *$f \in C(T, e)$  implies that  $T + e - f$  is a spanning tree of  $G$ .*



**Proof** (a) Lemma 2(a) implies that  $T + e$  has a cycle  $C$ . Suppose that  $T + e$  contains another cycle  $C' \neq C$ . Let edge  $g \in C' \setminus C$ .  $T' = T + e - g$  is connected, has  $n - 1$  edges. But  $T'$  contains a cycle  $C$ , contradicting Theorem 1.

(b)  $T + e - f$  is connected and has  $n - 1$  edges. Therefore it is a tree.  $\square$


## Maximum weight trees

$G = (V, E)$  is a connected graph.

$w : E \rightarrow \mathbf{R}$ .  $w(e)$  is the *weight* of edge  $e$ .

For spanning tree  $T$ ,  $w(T) = \sum_{e \in T} w(e)$ .

**Problem:** find a spanning tree of maximum weight.



## Greedy Algorithm

Sort edges so that  $E = \{e_1, e_2, \dots, e_m\}$  where

$$w(e_1) \geq w(e_2) \geq \dots \geq w(e_m).$$

begin

$T := \emptyset$

for  $i = 1, 2, \dots, m$  do

begin

if  $T + e_i$  does not contain a cycle

then  $T \leftarrow T + e_i$

end

Output  $T$

end

Greedy always adds the maximum weight edge which does not make a cycle with previously chosen edges.

**Theorem 4** *Let  $G$  be a connected weighted graph. The tree constructed by GREEDY is a maximum weight spanning tree.*

**Proof** Lemma 3 implies that  $T$  is a spanning tree of  $G$ .

Let the edges of the *greedy tree* be

$e_1^*, e_2^*, \dots, e_{n-1}^*$ , in order of choice. Note that  $w(e_i^*) \geq w(e_{i+1}^*)$  since neither makes a cycle with  $e_1^*, e_2^*, \dots, e_{i-1}^*$ .

Let  $f_1, f_2, \dots, f_{n-1}$  be the edges of any other tree where  $w(f_1) \geq w(f_2) \geq \dots \geq w(f_{n-1})$ .

We show that


$$w(e_i^*) \geq w(f_i) \quad 1 \leq i \leq n - 1. \quad (1)$$

Suppose (1) is false. There exists  $k > 0$  such that

$$w(e_i^*) \geq w(f_i), \quad 1 \leq i < k \text{ and } w(e_k^*) < w(f_k).$$

Each  $f_i$ ,  $1 \leq i \leq k$  is either one of or makes a cycle with  $e_1^*, e_2^*, \dots, e_{k-1}^*$ . Otherwise one of the  $f_i$  would have been chosen in preference to  $e_k^*$ .

Let components of forest  $(V, \{e_1^*, e_2^*, \dots, e_{k-1}^*\})$  be  $C_1, C_2, \dots, C_{n-k+1}$ . Each  $f_i$ ,  $1 \leq i \leq k$  has both of its endpoints in the same component.



Let  $\mu_i$  be the number of  $f_j$  which have both endpoints in  $C_i$  and let  $\nu_i$  be the number of vertices of  $C_i$ . Then

$$\mu_1 + \mu_2 + \cdots + \mu_{n-k+1} = k \quad (2)$$

$$\nu_1 + \nu_2 + \cdots + \nu_{n-k+1} = n \quad (3)$$

It follows from (2) and (3) that there exists  $t$  such that

$$\mu_t \geq \nu_t. \quad (4)$$

[Otherwise


$$\begin{aligned} \sum_{i=1}^{n-k+1} \mu_i &\leq \sum_{i=1}^{n-k+1} (\nu_i - 1) \\ &= \sum_{i=1}^{n-k+1} \nu_i - (n - k + 1) \\ &= k - 1. \end{aligned} \quad ]$$

But (4) implies that the edges  $f_j$  such that  $f_j \subseteq C_t$  contain a cycle.  $\square$

## Cut Sets and Bonds

If  $S \subseteq V$ ,  $S \neq \emptyset, V$  then the **cut-set**

$$S : \bar{S} = \{e = vw \in E : v \in S, w \in \bar{S} = V \setminus S\}$$



$$S = \{1, 2, 3\} \quad S : \bar{S} = \{d, e, f\}.$$



**Lemma 4** *Let  $G$  be connected and  $X \subseteq E$ . Then  $G[E \setminus X]$  is not connected iff  $X$  contains a cutset.*


**Proof**

**Only if**

$G[E \setminus X]$  contains components  $C_1, C_2, \dots, C_k$ ,  $k \geq 2$  and so  $X \supseteq C_1 : \bar{C}_1$  and  $C_1 \neq \emptyset, V$ .


**If**

Suppose  $X = S : \bar{S}$  and  $v \in S, w \in \bar{S}$ . Then every walk from  $v$  to  $w$  in  $G$  contains an edge of  $X$ .



So  $G[E \setminus X]$  contains no walk from  $v$  to  $w$ . □

A **Bond**  $B$  is a *minimal* cut-set. I.e.  $B = S : \bar{S}$  and if  $T : \bar{T} \subseteq B$  then  $B = T : \bar{T}$ .



$S_1 = \{1, 2, 3\}$      $B_1 = S_1 : \bar{S}_1$  is a bond

$S_2 = \{2, 3, 4, 5\}$      $B_2 = S_2 : \bar{S}_2$  is not a bond

since  $B_2 \supset S_3 : \bar{S}_3$  and  $B_2 \neq S_3 : \bar{S}_3$  where  $S_3 = \{1\}$ .

**Theorem 5**  $G$  is connected and  $B$  is a a bond  $\leftrightarrow G \setminus B$  contains exactly 2 components.

**Proof**  $\rightarrow$ :  $G \setminus B$  contains components  $C_1, C_2, \dots, C_k$ . Assume w.l.o.g. that there is at least one edge  $e$  in  $G$  joining  $C_1$  and  $C_2$ . If  $k \geq 3$  then  $B \supseteq C_3 : \bar{C}_3$  and  $B \neq C_3 : \bar{C}_3$  since  $B$  contains  $e$ .

$\leftarrow$ : Assume that  $G \setminus B$  contains exactly two components  $C_1 = G[S], C_2 = G[\bar{S}]$ . Let  $e \in B$ . Adding  $e$  to the graph  $C_1 \cup C_2$  clearly produces a connected graph and so  $B \setminus e$  is not a cutset.  $\square$

A **co-tree**  $\bar{T}$  of a connected graph  $G$  is the edge complement of a spanning tree of  $G$  i.e.  $\bar{T} = E \setminus T$  for some spanning tree  $T$ .

**Theorem 6** *Let  $T$  be a spanning tree of  $G$  and  $e \in T$ . Then*

**(a)**  $\bar{T}$  contains no bond of  $G$ .

**(b)**  $\bar{T} + e$  contains a unique bond  $B(\bar{T}, e)$  of  $G$ .

**(c)**  $f \in B(\bar{T}, e)$  implies that  $\bar{T} + e - f$  is a co-tree of  $G$ .


[Compare with Tree + edge  $\supseteq$  cycle.]

**Proof** (a)  $X \subseteq \bar{T} \leftrightarrow G \setminus X \supseteq T$  which implies that  $G \setminus X$  is connected. So  $X$  is not a bond.

(b)&(c)  $G \setminus (\bar{T} + e) = T \setminus e$  contains exactly two components and so by Theorem 5  $\bar{T} + e$  contains a bond  $B = S : \bar{S}$  where  $S, \bar{S}$  are the 2 components of  $T \setminus e$ .

$$\begin{aligned} f \in B &\Rightarrow e \in C(T, f) \\ &\Rightarrow T + f - e \text{ is a tree} \\ &\Rightarrow \bar{T} + e - f \text{ is a co-tree} \quad \text{proving (c)} \end{aligned}$$


Hence every bond of  $\bar{T} + e$  contains  $f$  – otherwise  $\bar{T} + e - f$  contains a bond, contradicting (a) and proving (b).  $\square$



$$B = (S : \bar{S})$$

# How many trees? – Cayley's Formula

n=4




4



12

n=5



5




60




60

n=6




6



120



360



90




360



360

## Contracting edges

If  $e = vw \in E$ ,  $v \neq w$  then we can **contract**  $e$  to get  $G \cdot e$  by (i) deleting  $e$ , (ii) identifying  $v, w$  i.e. make them into a single new vertex.



$G - e$  is obtained by deleting  $e$ .

$\tau(G)$  is the number of spanning trees of  $G$ .



**Theorem 7** *If  $e \in E$  is not a loop then*

$$\tau(G) = \tau(G \cdot e) + \tau(G - e).$$


**Proof**

- $\tau(G - e)$  = the number of spanning trees of  $G$  which do not contain  $e$ .
- $\tau(G \cdot e)$  = the number of spanning trees of  $G$  which contain  $e$ .

[Bijection  $T \rightarrow T \cdot e$  maps spanning trees of  $G$  which contain  $e$  to spanning trees of  $G \cdot e$ .]

## Matrix Tree Theorem

Define the  $V \times V$  matrix  $L = D - A$  where  $A$  is the adjacency matrix of  $G$  and  $D$  is the diagonal matrix with  $D(v, v) = \text{degree of } v$ .



$$L = \begin{vmatrix} 3 & -1 & -1 & -1 \\ -1 & 3 & -1 & -1 \\ -1 & -1 & 3 & -1 \\ -1 & -1 & -1 & 3 \end{vmatrix}$$

$$L_1 = \begin{vmatrix} 3 & -1 & -1 \\ -1 & 3 & -1 \\ -1 & -1 & 3 \end{vmatrix}$$

$$\text{Determinant } L_1 = 16$$

Let  $L_1$  be obtained by deleting the first row and column of  $L$ .

### Theorem 8

$$\tau(G) = \text{determinant } L_1.$$

## Pfuffer's Correspondence

There is a 1-1 correspondence  $\phi_V$  between spanning trees of  $K_V$  (the complete graph with vertex set  $V$ ) and sequences  $V^{n-2}$ . Thus for  $n \geq 2$

$$\tau(K_n) = n^{n-2} \quad \text{Cayley's Formula.}$$

Assume some arbitrary ordering  $V = \{v_1 < v_2 < \dots < v_n\}$ .

$\phi_V(T)$ :

**begin**

$T_1 := T;$

**for**  $i = 1$  **to**  $n - 2$  **do**


**begin**

$s_i :=$  neighbour of least leaf  $\ell_i$  of  $T_i$ .

$T_{i+1} = T_i - \ell_i$ .

**end**  $\phi_V(T) = s_1 s_2 \dots s_{n-2}$

**end**




6,4,5,14,2,6,11,14,8,5,11,4,2

**Lemma 5**  $v \in V(T)$  appears exactly  $d_T(v) - 1$  times in  $\phi_V(T)$ .

**Proof** Assume  $n = |V(T)| \geq 2$ . By induction on  $n$ .

$n = 2$ :  $\phi_V(T) = \Lambda =$  empty string.

Assume  $n \geq 3$ :



$\phi_V(T) = s_1 \phi_{V_1}(T_1)$  where  $V_1 = V - \{l_1\}$ .

$s_1$  appears  $d_{T_1}(s_1) - 1 + 1 = d_T(s_1) - 1$  times – induction.

$v \neq s_1$  appears  $d_{T_1}(v) - 1 = d_T(v) - 1$  times – induction.  $\square$

## Construction of $\phi_V^{-1}$

Inductively assume that for all  $|X| < n$  there is an inverse function  $\phi_X^{-1}$ . (True for  $n = 2$ ).

Now define  $\phi_V^{-1}$  by

$\phi_V^{-1}(s_1 s_2 \dots s_{n-2}) = \phi_{V_1}^{-1}(s_2 \dots s_{n-2})$  plus edge  $s_1 \ell_1$ ,

where  $\ell_1 = \min\{s : s \notin s_1, s_2, \dots, s_{n-2}\}$  and  $V_1 = V - \{\ell_1\}$ .

Then

$$\begin{aligned}\phi_V(\phi_V^{-1}(s_1 s_2 \dots s_{n-2})) &= \\ &= \phi_V(\phi_{V_1}^{-1}(s_2 \dots s_{n-2}) \text{ plus edge } s_1 \ell_1) \\ &= s_1 \phi_{V_1}(\phi_{V_1}^{-1}(s_2 \dots s_{n-2})) \\ &= s_1 s_2 \dots s_{n-2}.\end{aligned}$$

Thus  $\phi_V$  has an inverse and the correspondence is established.

## Number of trees with a given degree sequence

**Corollary 5** *If  $d_1 + d_2 + \cdots + d_n = 2n - 2$  then the number of spanning trees of  $K_n$  with degree sequence  $d_1, d_2, \dots, d_n$  is*

$$\binom{n-2}{d_1-1, d_2-1, \dots, d_n-1} = \frac{(n-2)!}{(d_1-1)!(d_2-1)! \cdots (d_n-1)!}$$

**Proof** From Puffer's correspondence and Lemma 5 this is the number of sequences of length  $n - 2$  in which 1 appears  $d_1 - 1$  times, 2 appears  $d_2 - 1$  times and so on. □