
NOTES ON GROWING A TREE IN A GRAPH^⓪

Luc Devroye,[§] Vida Dujmović,[¶] Alan Frieze,[♠] Abbas Mehrabian,^⓪ Pat Morin,^² and Bruce Reed[♣]

ABSTRACT. We study the height of a spanning tree T of a graph G obtained by starting with a single vertex of G and repeatedly selecting, uniformly at random, an edge of G with exactly one endpoint in T and adding this edge to T .

^⓪This research was partly supported by NSERC and by NSF GRANT DMS1661063. Some of this work was carried out while Vida Dujmović, Pat Morin, and Bruce Reed were visiting IMPA in Rio de Janeiro. They thank the Institute for its hospitality.

[§]School of Computer Science, McGill University, Montréal, Canada

[¶]Department of Computer Science and Electrical Engineering, University of Ottawa, Ottawa, Canada

[♠]Department of Mathematical Sciences, Carnegie Mellon University, Pittsburgh, USA. Research supported in part by NSF grant DMS1362785.

^⓪Department of Computer Science, University of British Columbia, Vancouver, Canada. This author was supported by an NSERC Postdoctoral Fellowship and a Simons-Berkeley Research Fellowship. Part of this work was done while he was visiting the Simons Institute for the Theory of Computing at UC Berkeley.

^²School of Computer Science, Carleton University, Ottawa, Canada

[♣]CNRS, Projet COATI, I3S (CNRS and UNS) UMR7271 and INRIA, Sophia Antipolis, France. Instituto Nacional de Matemática Pura e Aplicada (IMPA), Brasil. Visiting Research Professor, ERATO Kawarabayashi Large Graph Project, Japan.

Contents

1	Introduction	1
2	Preliminaries	2
3	Connection with first-passage percolation and a generic upper bound	3
4	An upper bound in terms of graph degeneracy	5
5	An upper bound in terms of Euler genus	7
6	An upper bound for edge expanders	8
7	Upper bounds for high dimensional grids and hypercubes	10
8	Lower Bounds for General Graphs	13
8.1	The Ladder Graph H	14
8.2	The Subdivided Tree I	15
8.3	Putting it Together	15
9	Lower Bounds for Degenerate Graphs	17
A	Proof of Lemma 1	22

1 Introduction

Let s be a vertex of a simple connected graph G on n vertices. We build a sequence T_1, T_2, \dots, T_n of random subtrees of G as follows. The tree T_1 has a single vertex, s . For each $1 < i \leq n$, tree T_i is obtained by choosing a uniformly random edge of G with exactly one endpoint in T_{i-1} , and adding the edge to T_{i-1} . Note that T_n is a (not necessarily uniform) random spanning tree of G rooted at s , which we denote by $\mathcal{T}(G, s)$. In this paper we study the height (maximum length of a root-to-leaf path) of $\mathcal{T}(G, s)$ and give several bounds for it in terms of parameters of G .

In the special case when G is the complete graph, each tree T_i is obtained from T_{i-1} by choosing a uniformly random node of T_{i-1} and joining a new leaf to that node. This is the well studied *random recursive tree* process, and Devroye [6] and Pittel [13] have shown that the height of $T_n = \mathcal{T}(K_n, s)$ is $(e + o(1)) \ln n$ with probability $1 - o(1)$.

Our results. Let $D = D(G)$ and $\Delta = \Delta(G)$ denote the diameter and maximum degree of G , respectively, and let us denote the height of a tree T by $h(T)$. An obvious lower bound for $h(\mathcal{T}(G, s))$ is $D/2$. We prove the following bounds hold with probability $1 - o_n(1)$ for any n -vertex graph G and any $s \in V(G)$. (The notation $o_k(1)$ denotes the set of functions $f : \mathbb{R} \rightarrow \mathbb{R}$ such that $f(k) \rightarrow 0$ as $k \rightarrow \infty$.)

1. In Theorem 1 we show $h(\mathcal{T}(G, s)) \in O(\Delta(D + \log n))$. For $D \in \Omega(\log \Delta)$ this is tight: in Theorem 7 we show that for every $\Delta \geq 2$ and every $D \geq e^6 \ln \Delta$, there exist G and s with $h(\mathcal{T}(G, s)) \in \Omega(\Delta(D + \log n))$.
2. If G is d -degenerate (that is, every subgraph of G has a vertex of degree at most d), then in Theorem 2 we show $h(\mathcal{T}(G, s)) \in O(\sqrt{d\Delta}(D + \log n))$. The class of $O(1)$ -degenerate graphs is quite rich and includes every minor-closed graph family. This upper bound is tight, even for planar graphs ($d = 5$), graphs of thickness t ($d = 5t$), and graphs of treewidth k ($d = k$). (The concepts of Euler genus, thickness, and treewidth are defined in Section 4).

For $D \in \Omega(\log \Delta)$ and planar graphs (which are 5-degenerate) this is tight: in Theorem 8 we show for any $\Delta > 2$ and $D > 10^6 \ln \Delta$ there exists a planar graph G and vertex s with $h(\mathcal{T}(G, s)) \in \Omega(\sqrt{\Delta}(D + \log n))$.

Also, for $D \in \Omega(\log \Delta)$ and $d \leq \Delta$ this is tight: in Theorem 9 we show for any $\Delta > 1$, $D > 10^6 \ln \Delta$ and $d \leq \Delta$ there exist a d -degenerate graph G and vertex s with $h(\mathcal{T}(G, s)) \in \Omega(\sqrt{d\Delta}(D + \log n))$.

3. If G has Euler genus less than $C\sqrt{\Delta}D/\log \Delta$, then $h(\mathcal{T}(G, s)) \in O(\sqrt{\Delta}(D + \log n))$ (see Theorem 3). For $D \in \Omega(\log \Delta)$ and zero Euler genus this is tight: in Theorem 8 we show for any $\Delta > 2$ and $D > 10^6 \ln \Delta$ there exist a planar graph G and vertex s with $h(\mathcal{T}(G, s)) \in \Omega(\sqrt{\Delta}(D + \log n))$.
4. For any $d, k \geq 1$, if G is the d -dimensional grid of side-length k (which has $n = (k+1)^d$ vertices), we have $h(\mathcal{T}(G, s)) \in O(dk + d^{5/3} \ln(k+1))$. If $k = 2$ or $k/\ln(k+1) = \Omega(d^{2/3})$, we have $h(\mathcal{T}(G, s)) \in \Theta(D) = \Theta(dk)$ (see Theorem 5 and Corollary 2).

-
5. If G has edge-expansion factor¹ (i.e., Cheeger constant) Φ , then $h(\mathcal{T}(G, s)) \in O(\Phi^{-1} \Delta \log n)$ (see Theorem 4). This implies, for example, that $h(\mathcal{T}(G, s)) \in O(\log n)$ if G is the complete graph or if G is a random Δ -regular graph (since a random Δ -regular graph has $\Phi \in \Omega(\Delta)$, see [3]).

Our main tool for proving upper bounds, Lemma 2, bounds $h(\mathcal{T}(G, s))$ in terms of the first-passage percolation cover time and the number of paths of a given length starting at s . To prove our results using this tool, we prove several new bounds on first-passage percolation cover times as well as the number of simple paths in various families of graphs, which are of independent interest.

Our results on first-passage percolation cover time. Suppose independent exponential(1) random variables $\{\tau_e\}$ are assigned to edges of G . Let $\Gamma(s, v)$ denote the set of all (s, v) -paths in the graph. Then the *first-passage percolation cover time* is defined as

$$\tau(G, s) = \max_{v \in V(G)} \min_{\gamma \in \Gamma(s, v)} \sum_{e \in \gamma} \tau(e)$$

In Lemma 3 we show a general upper bound of $O(\ln n + D)$ for $\tau(G, s)$. (This and the following results hold with probability $1 - o_n(1)$.)

In the special case when G is the d -dimensional grid with side length k (and diameter dk), we prove the improved bound $\tau(G, s) = O(k)$. The special case of $k = 1$, namely the d -cube graph, was studied by Fill and Pemantle [8], who showed $1.414 \leq \tau(G, s) \leq 14.041$. The upper bound was subsequently improved to 1.694 by Bollobás and Kohayakawa [4] and recently to 1.575 by Martinsson [10].

The remainder of the paper is organized as follows: Section 2 presents some preliminaries and useful facts about sums of independent random variables, In Section 3 we present the connection with first-passage percolation and prove a general upper bound. Section 4–Section 7 present our upper bounds on $h(\mathcal{T})$. Section 8 and Section 9 present families of graphs with matching lower bounds.

We use the following notational conventions: $\log x$ denotes the binary logarithm of x and $\ln x$ denotes the natural logarithm of x . Every graph, G , that we consider is finite, simple, undirected and connected, and n denotes its number of vertices.

2 Preliminaries

Recall that an exponential(λ) random variable, X , has a distribution defined by

$$\Pr\{X > x\} = e^{-\lambda x}, \quad x \geq 0,$$

and mean $E[X] = \int_0^\infty \Pr\{X > x\} dx = 1/\lambda$. We make extensive use of the fact that exponential random variables are *memoryless*:

$$\Pr\{X > t + x \mid X > t\} = \frac{\Pr\{X > t + x\}}{\Pr\{X > t\}} = \frac{e^{-\lambda(t+x)}}{e^{-\lambda t}} = e^{-\lambda x} = \Pr\{X > x\}.$$

¹The edge expansion factor and related quantities are defined in Section 6.

We will also often take the minimum of δ independent $\text{exponential}(\lambda)$ random variables and use the fact that this is distributed like an $\text{exponential}(\lambda\delta)$ random variable:

$$\Pr\{\min\{X_1, \dots, X_\delta\} > x\} = (\Pr\{X_1 > x\})^\delta = e^{-\delta\lambda x} \sim \text{exponential}(\lambda\delta) .$$

We will make use of two concentration inequalities for sums of exponential random variables, both of which can be obtained using Chernoff's bounding method (see, e.g., [9, Theorem 5.1]). If Z_1, \dots, Z_k are independent $\text{exponential}(\lambda)$ random variables (so that they each have mean $\mu = 1/\lambda$), then for all $d > 1$,

$$\Pr\left\{\sum_{i=1}^k Z_i \leq \mu k/d\right\} \leq \exp(-k(\ln d - 1 + 1/d)) \leq \left(\frac{e}{d}\right)^k \quad (1)$$

and for all $t > 1$,

$$\Pr\left\{\sum_{i=1}^k Z_i \geq \mu kt\right\} \leq \exp(k - kt/2) . \quad (2)$$

The sum of k independent $\text{exponential}(\lambda)$ random variables is called an $\text{Erlang}(k, \lambda)$ random variable.

For positive integers a and b , we define the random variable $Y_{a,b}$ as follows: Consider a tree in which the root has a children, and each of the root's children have b children. Put an independent $\text{exponential}(1)$ weight on each edge. Then $Y_{a,b}$ is defined as the minimum weight of a path from the root to a leaf. The following auxiliary lemma is proved in Appendix A.

Lemma 1. *Let X_1, \dots, X_m be i.i.d. distributed as $Y_{a,b}$ for some a, b . Then*

$$\mathbb{E}[X_1] = O(1/a + 1/\sqrt{ab})$$

and moreover,

$$\Pr\left\{\sum_{i=1}^m X_i \geq 3m(64/a + 1024/\sqrt{ab})\right\} \leq \exp(-m/9).$$

3 Connection with first-passage percolation and a generic upper bound

In this section, we establish the connection with first-passage percolation, and prove an upper bound for $\tau(G, s)$ in general graphs, which results in an upper bound for $h(\mathcal{T}(G, s))$. This connection will be used in subsequent sections to provide tighter bounds for $h(\mathcal{T}(G, s))$ in several graph classes.

Recall the generation process for $\mathcal{T}(G, s)$: we start with a tree containing only vertex s initially; in each round, we choose an edge uniformly at random among edges with exactly one endpoint in the existing tree, and add it to the existing tree.

We may view this as an *infection process*: at round 0 only vertex s is infected. In each round, suppose the set of infected vertices is S . We choose a uniformly random edge

between S and its complement, and let the disease spread along that edge, hence increasing the number of infected vertices by one.

Now consider the following continuous time view of this infection process, which is known as Richardson's model [7] or first-passage percolation [1]. At time 0 we infect vertex s . For each edge uv , whenever one of u and v gets infected, we put an exponential(1) timer on edge uv . When the timer rings, the disease spreads along that edge and both u and v get infected (it might be the case that both u and v are already infected by that time). Suppose at some moment in this process, the subset S of vertices are infected. Then, by memorylessness of the exponential distribution, the disease is equally like to spread along any of the edges existing between S and its complement. Therefore, the tree along which the disease spreads has the same distribution as $\mathcal{T}(G, s)$.

This viewpoint induces weights on the edges: to each edge e we assign weight $\tau(e)$, which is the ringing time for the timer on this edge. Note that the weights are i.i.d. exponential (1) random variables. The weight of a path P , denoted $\tau(P)$, is simply the sum of weights of its edges. The *first-passage percolation hitting time* (or simply, the *hitting time*) for v is the weight of the lightest path from s to v :

$$\tau(G, s, v) = \min_{\gamma \in \Gamma(s, v)} \tau(\gamma) .$$

The *first-passage percolation cover time* (or simply, the *cover time*) is the first time that all vertices are infected, which can be written as

$$\tau(G, s) = \max_{v \in V(G)} \tau(G, s, v) .$$

Note that this is also the maximum *weight* of a root-to-leaf path in the infection tree $\mathcal{T}(G, s)$, which we will use to bound the height of $\mathcal{T}(G, s)$, the maximum *length* of a path in (the unweighted version of) $\mathcal{T}(G, s)$ (in general, the longest path and the heaviest path may be different).

For a positive integer L and a vertex s of graph G , let $\Pi(G, s, L)$ denote the number of simple paths of length L in G that start from s . We now prove a lemma that upper bounds $h(\mathcal{T}(G, s))$ in terms of $\tau(G, s)$ and $\Pi(G, s, L)$.

Lemma 2. *Let $s \in V(G)$, $0 \leq p < 1 \leq a$, $c > 0$, and $L = \lceil ceaK \rceil$ be such that $\Pr\{\tau(G, s) > K\} \leq p$ and $\Pi(G, s, L) \leq a^L$. Then $h(\mathcal{T}(G, s)) \leq L$ with probability at least $1 - p - c^{-L}$.*

Proof. Let $T = \mathcal{T}(G, s)$. If $h(T) > L$, then at least one of the following two events occurred:

1. T contains a root-to-leaf path of weight greater than K .
2. G contains a path starting at s of length L whose weight is less than K .

By assumption, the probability of the first event is at most p . The weight of a single path of length L is the sum of L exponential(1) random variables so, by (1) and the union bound over all a^L paths, the probability of the second event is at most

$$a^L \left(\frac{eK}{L} \right)^L \leq c^{-L} . \quad \square$$

In light of Lemma 2, we can upper bound $h(\mathcal{T}(G, s))$ if we have upper bounds on the cover time and on the number of paths of length L originating at s . An obvious upper bound for the latter is Δ^L . The following lemma gives a general upper bound for the former, which results in a general upper bound for $h(\mathcal{T}(G, s))$. In the following sections we obtain better bounds for these two quantities in special graph classes, resulting in sharper bounds on $h(\mathcal{T}(G, s))$.

Lemma 3. *For any $s \in V(G)$, we have $\tau(G, s) \leq 4\ln n + 2D$ with probability at least $1 - 1/n$.*

Proof. For each vertex $v \in V(G)$, we show the probability that it is not infected by time $4\ln n + 2D$ is at most n^{-2} , and then apply the union bound over all vertices. Let P be a shortest (s, v) -path in G . Let $k \leq D$ denote the length of P , so $\tau(P) \sim \text{Erlang}(k, 1)$. Note that for any t , $\tau(P) \leq t$ implies v is infected by time t . Thus, using (2), the probability that v is not infected by time $4\ln n + 2D$ is bounded by

$$\Pr\{\tau(P) > 4\ln n + 2D\} = \Pr\{\text{Erlang}(k, 1) > 4\ln n + 2D\} \leq \exp(k - 2\ln n - D) \leq n^{-2}. \quad \square$$

We immediately get a general upper bound for $h(\mathcal{T}(G, s))$.

Theorem 1. *Let G be an n -vertex graph with diameter D and maximum degree $\Delta > 1$, and let s be an arbitrary vertex. Then, with probability at least $1 - O(1/n)$ we have*

$$\frac{D}{2} \leq h(\mathcal{T}(G, s)) \leq 2e\Delta(4\ln n + 2D) \leq (4e\Delta + 8e\Delta \ln \Delta)D + 16e\Delta.$$

Note that this gives an asymptotically tight bound of $h(\mathcal{T}(G, s)) = \Theta(D)$ for graphs with bounded maximum degree.

Proof. The first inequality is trivial. The second inequality is an application of Lemma 2 with $a = \Delta$, $p = 1/n$, $K = 4\ln n + 2D$ and $c = 2$, using the bound of Lemma 3 for the cover time. The last inequality follows from the crude bound $\Delta^D \geq n/3$, which holds for any n -vertex graph with maximum degree Δ and diameter D . \square

4 An upper bound in terms of graph degeneracy

Recall that a graph is d -degenerate if each of its subgraphs has a vertex of degree at most d . The following lemma shows that, for large L , d -degenerate graphs have considerably less than Δ^L walks of length L .

Lemma 4. *Let G be an n -vertex d -degenerate graph with maximum degree Δ . Then the number of walks in G of length L is bounded by $2n2^L(d\Delta)^{L/2}$.*

Proof. Enumerate the vertices of G as v_1, \dots, v_n so that v_i has at most d edges in the subgraph induced by v_i, \dots, v_n (this ordering may be obtained by repeatedly removing a vertex of degree at most d).

We give a way to encode the walks in a one-to-one way, and then bound the total number of possible generated codes. Let $W = v_{i_0}, \dots, v_{i_L}$ be a walk of length L in G and let $k = k(W)$ denote the number of indices $\ell \in \{1, \dots, L\}$ such that $i_{\ell-1} < i_\ell$. If $k \geq L/2$ then we say that W is *easy*; note that at least one of W and its reverse is easy, hence the total number of L -walks is at most twice the number of easy L -walks. We encode an easy walk W in the following way:

1. We first specify the starting vertex v_{i_0} . There are n ways to do this.
2. Next we specify whether $i_{\ell-1} < i_\ell$ for each $\ell \in \{1, \dots, L\}$. There are at most 2^L ways to do this.
3. Next, we specify each edge of W . For each $\ell \in \{1, \dots, L-1\}$, if $i_\ell < i_{\ell+1}$, then there are at most d ways to do this, otherwise there are at most Δ ways to do this. Therefore, the total number of ways to specify all edges of the walk is at most

$$d^k \Delta^{L-k} \leq (d\Delta)^{L/2} ,$$

since $d \leq \Delta$ and $k \geq L/2$.

Therefore, the number of easy L -walks is bounded by $n2^L(d\Delta)^{L/2}$, as required. \square

Theorem 2. *Let G be an n -vertex d -degenerate graph with diameter D and maximum degree Δ , and let s be an arbitrary vertex. Then, with probability at least $1 - O(1/n)$ we have $h(\mathcal{T}(G, s)) \leq 8e\sqrt{d\Delta}(2D + 4\ln n)$.*

Proof. Let $c = 2$, $K = 4\ln n + 2D$, $p = 1/n$, $a = 4\sqrt{d\Delta}$, and $L = \lceil ceaK \rceil > 8\ln n$. Lemma 3 guarantees $\tau(G, s) \leq 4\ln n + 2D$ with probability at least $1 - 1/n$, and Lemma 4 guarantees $\Pi(G, s, L) \leq 2n2^L(d\Delta)^{L/2} \leq a^L$. Applying Lemma 2 completes the proof. \square

Note that Theorem 2 actually implies Theorem 1 up to constant factors, since all graphs of maximum degree Δ are Δ -degenerate, so $\sqrt{d\Delta} \leq \Delta$ in all cases. However, Theorem 2 provides sharper bounds for many important graph classes:

- Planar graphs are 5-degenerate. (This is a consequence of Euler's formula and the fact that planarity is preserved under taking subgraphs).
- The *thickness* of a graph is the minimum number of planar graphs into which the edges of G can be partitioned. Graphs of thickness t are $5t$ -degenerate. (This follows from definitions and the 5-degeneracy of each individual planar graph in the partition.)
- The *Euler genus* of a graph is the minimum Euler genus of a surface on which the graph can be drawn without crossing edges. Graphs of Euler genus g are $O(\sqrt{g})$ -degenerate.²

²This follows from the facts in every n -vertex Euler-genus g graph, $n \in \Omega(\sqrt{g})$ and there exists a vertex of degree at most $6 + O(g/n)$. (See, e.g., [14, Lemma 7 and Theorem 2].)

-
- A *tree decomposition* of a graph G is a tree T' whose vertex set B is a collection of subsets of $V(G)$ called *bags* with the following properties:

1. For each edge vw of G , there is at least one bag $b \in B$ with $\{v, w\} \subseteq B$.
2. For each a vertex v of G , the subgraph of T' induced by the set of bags that contain v is connected.

The *width* of a tree-decomposition is one less than the size of its largest bag. The *treewidth* of G is the minimum width of any tree decomposition of G . Graphs of treewidth k are k -degenerate. (This is a consequence of the fact that k -trees are edge-maximal graphs of treewidth k .)

Therefore, Theorem 2 implies that, when the relevant parameter, g , t or k , is bounded, $h(T) \in O(\sqrt{\Delta}(D + \log n))$ with high probability.

5 An upper bound in terms of Euler genus

Since graphs of Euler genus g are $O(\sqrt{g})$ -degenerate, Theorem 2 implies that if G has Euler genus g , then $h(\mathcal{T}(G, s)) \in O(g^{1/4}\Delta^{1/2}(D + \log n))$. In this section we show that the dependence on the genus g can be eliminated when the diameter is large compared to the genus. We begin with an upper-bound on path counts that is better (for graphs of small genus) than Lemma 4.

Lemma 5. *Let G be a simple n -vertex graph of Euler genus g , diameter D , and maximum degree $\Delta \geq 6$. Then the number of simple paths in G of length L is at most $2n2^L 6^{L/2-3g} \Delta^{L/2+3g}$.*

Proof. The following proof makes use of some basic notions related to graphs on surfaces; see Mohar and Thomassen [12] for basic definitions and results. Since G has Euler genus g , it has a 2-cell embedding in a surface of Euler genus g . Euler's formula then states that

$$m = n + f - 2 + g, \quad (3)$$

where n and m are the numbers of vertices and edges of G and f is the number of faces in the embedding of G . Every edge is on the boundary of at most 2 faces of the embedding and, since G is simple, every face is bounded by at least 3 edges. Therefore, $f \leq 2m/3$, so (3) implies

$$m \leq 3n - 6 + 3g.$$

Therefore, the average degree of an n -vertex Euler genus g graph is at most $6 + (6g - 12)/n$. In particular, if $n \geq 6g$, then g has average degree less than 7, so G contains a vertex of degree at most 6.

When we remove a vertex from G we obtain a graph whose Euler genus is not more than that of G . Therefore, by repeatedly removing a degree 6 vertex, we can order the vertices of G as v_1, \dots, v_n so that, for each $i \in \{1, \dots, n - 6g\}$, v_i has at most 6 neighbours among v_{i+1}, \dots, v_n . We call v_{n-6g+1}, \dots, v_n *annoying vertices* and edges between them are *annoying edges*.

Let $P = v_{i_0}, \dots, v_{i_L}$ be a path of length L in G . For each $i \in \{1, \dots, L\}$, the edge $v_{i_{\ell-1}}v_{i_\ell}$ is called *bad* if it is annoying or if $i_{\ell-1} > i_\ell$; otherwise it is called good. Let k denote the number of good edges in P . Say P is good if $k \geq L/2 - 3g$. Note that the number of annoying edges of P is bounded by $6g - 1$, hence at least one of P and its reverse is good. We bound the number of good L -paths; the total number of L -paths is at most twice this bound. We encode a good L -path P as follows:

1. We first specify the starting vertex v_{i_0} . There are n ways to do this.
2. Next we specify whether each edge of P is good or bad. There are 2^L ways to do this.
3. Next, we specify each edge of P . For each good edge, there are at most 6 ways to do this. For each bad edge there are at most Δ ways to do this. Therefore, the total number of ways to specify the edges of P is at most

$$6^k \Delta^{L-k} \leq 6^{L/2-3g} \Delta^{L/2+3g} ,$$

since $k \geq L/2 - 3g$ and $\Delta \geq 6$.

Therefore, the number of good L -paths is at most $n 2^L 6^{L/2-3g} \Delta^{L/2+3g}$, as required. \square

Theorem 3. *Let G be an n -vertex Euler-genus g graph with diameter D , maximum degree Δ and let $s \in V(G)$ be an arbitrary vertex. If $g \ln \Delta \leq 36\sqrt{\Delta}(D + \ln n)$ then, with probability at least $1 - O(1/n)$, $h(T(G, s)) \leq 107\sqrt{\Delta}(2D + 4\ln n)$.*

Proof. The conclusion follows from Theorem 1 for $\Delta \leq 6$, so we will assume $\Delta > 6$. Let $c = 2$, $K = 4\ln n + 2D$, $p = 1/n$, $a = 8\sqrt{6\Delta}$, and $L = \lceil ceaK \rceil > 8\ln n$. Lemma 3 guarantees $\tau(G, s) \leq 4\ln n + 2D$ with probability at least $1 - 1/n$, and Lemma 4 guarantees

$$\Pi(G, s, L) \leq 2n \times 2^L \times (6\Delta)^{L/2} \times \Delta^{3g} \leq (2 \times 2 \times \sqrt{6\Delta})^L \exp(108\sqrt{\Delta}(D + \ln n)) \leq (2 \times 2 \times \sqrt{6\Delta} \times 2)^L = a^L.$$

Applying Lemma 2 completes the proof. \square

6 An upper bound for edge expanders

All of the preceding upper bounds on $h(T)$ have a (linear or rootish) dependence on Δ , the maximum degree of a vertex in G . This seems somewhat counterintuitive, since high degree vertices in G should produce high degree vertices in T and therefore decrease $h(T)$. In this section we show that indeed large edge expansion (also called isoperimetric number or Cheeger constant) results in low-height trees.

For an n -vertex graph G and a subset $A \subseteq V(G)$, define $e(A) = |\{vw \in E(G) : v \in A, w \notin A\}|$, and for any $k \in \{1, \dots, n-1\}$, define

$$e_k(G) = \min\{e(A) : A \subseteq V(G), |A| = k\} .$$

Observe that $e_k(G)$ is symmetric in the sense that $e_k(G) = e_{n-k}(G)$. We define the *edge expansion* of G is

$$\Phi(G) = \min\{e_k(G)/k : k \in \{1, \dots, \lfloor n/2 \rfloor\}\}$$

We will express the height of T in terms of the *total inverse perimeter size* Ψ , which is closely related to the edge expansion:

$$\Psi(G) = \sum_{k=1}^{\lfloor n/2 \rfloor} \frac{1}{e_k(G)} \leq \sum_{k=1}^{\lfloor n/2 \rfloor} \frac{1}{k\Phi(G)} = \frac{\ln n + O(1)}{\Phi(G)}.$$

Theorem 4. *Let G be an n -vertex graph with maximum degree Δ , edge-expansion Φ , total inverse perimeter size Ψ , and let s be an arbitrary vertex. Then, with probability at least $1 - \exp(-\Omega(\Psi\Delta))$ we have $h(\mathcal{T}(G, s)) \in O(\Psi\Delta) \subseteq O(\Phi^{-1}\Delta \log n)$.*

Before proving Theorem 4, we consider the example of the complete graph $G = K_n$. In this graph, the minimum degree is $n - 1$, so all preceding theorems (at best) imply an upper bound of $O(n)$ on $h(\mathcal{T}(K_n, s))$. However, $e_k(K_n) = k(n - k)$, so $\Phi(K_n) = \lceil n/2 \rceil$, and $\Psi(K_n) = O(\log n/n)$. Then Theorem 4 implies that $h(\mathcal{T}(K_n, s)) \in O(\log n)$ with high probability. This upper bound is of the right order of magnitude, since it matches the (tight) results of Devroye and Pittel for the height of the random recursive tree [6, 13].

Proof. Fix some path $P = (s = v_0), v_1, \dots, v_L$ in G and suppose that P appears as a path in T . Then there are times $1 \leq k_1 < \dots < k_L < n$ such that for each $i \in \{1, \dots, L\}$, v_i joins T when T has size k_i . For a fixed P and fixed $1 \leq k_1 < \dots < k_L < n$, the probability that this happens is at most

$$\prod_{i=1}^L \frac{1}{e_{k_i}(G)},$$

and the probability that P appears in T (without fixing k_1, \dots, k_L) is at most

$$\sum_{1 \leq k_1 < \dots < k_L < n} \left(\prod_{i=1}^L \frac{1}{e_{k_i}(G)} \right) < \frac{1}{L!} \left(\sum_{(k_1, \dots, k_L) \in \{1, \dots, n-1\}^L} \left(\prod_{i=1}^L \frac{1}{e_{k_i}(G)} \right) \right) = \frac{1}{L!} \left(\sum_{k=1}^{n-1} \frac{1}{e_k(G)} \right)^L \leq \frac{(2\Psi)^L}{L!}$$

Finally, since G contains at most Δ^L paths of length L ,

$$\Pr\{h(\mathcal{T}(G, s)) \geq L\} \leq \Delta^L \times \frac{(2\Psi)^L}{L!} \leq \left(\frac{2e\Psi\Delta}{L} \right)^L \leq \left(\frac{1}{2} \right)^L,$$

for $L \geq 4e\Psi\Delta$. □

Observe that the last step in the proof of Theorem 4 is to use the union bound over all paths of length L . If we have a better upper-bound than Δ^L on the number of such paths, then we obtain a better upper bound on $h(T)$. Lemma 4 gives a better upper bound for d -degenerate graphs, using which we immediately obtain the following corollary.

Corollary 1. *Let G be an n -vertex d -degenerate graph with diameter D and maximum degree Δ , total inverse perimeter size Ψ , and let s be an arbitrary vertex. Then, with probability at least $1 - O(1/n)$, $h(\mathcal{T}(G, s)) \in O(\Psi\sqrt{d\Delta} + \log n) \in O(\log n(1 + \sqrt{d\Delta}/\Phi))$.*

Proof. As in the proof of Theorem 4, and using the upper bound $2n2^L(d\Delta)^{L/2}$ for the number of paths of length L , given by Lemma 4, we have

$$\Pr\{h(\mathcal{T}(G,s)) \geq L\} \leq 2n2^L(d\Delta)^{L/2} \times \frac{(2\Psi)^L}{L!} \leq 2n(4e\Psi\sqrt{d\Delta}/L)^L \leq (8e\Psi\sqrt{d\Delta}n^{1/L}/L)^L,$$

which is smaller than $1/n$ for $L \geq 8e^3\Psi\sqrt{d\Delta} + \ln n$, as required. \square

7 Upper bounds for high dimensional grids and hypercubes

The d -cube is the graph having vertex set $\{0,1\}^d$ in which two vertices are adjacent if and only if they differ in exactly one coordinate. Every vertex in the d -cube has degree d and the d -cube has diameter d . The d -cube is an interesting example in which the path count is high, but this is counteracted by a low first-passage percolation time.

Theorem 5. *Let $n = 2^d$, let G be the d -cube and let $s \in V(G)$ be arbitrary. Then, with probability at least $1 - o_n(1)$, $h(\mathcal{T}(G,s)) \in \Theta(d)$.*

Proof. Fill and Pemantle [8] showed that the first-passage percolation cover time for the d -cube is at most 14.05 with probability $1 - o_n(1)$. Every vertex of the hypercube has degree d , so the number of paths of length L starting at s is less than d^L . The result then follows by applying Lemma 2 with $p = o_n(1)$, $c = 2$, $K = 14.05$, and $a = d$. \square

A natural generalization of the d -cube is the (d,k) -grid, which has vertex set $\{0, \dots, k\}^d$ and has an edge between two vertices if and only if the (Euclidean or ℓ_1) distance between them is 1. The (d,k) -grid has diameter $D = dk$ and maximum degree $\Delta = 2d$.

Note that in the case $k = 1$, the $(d,1)$ -grid is the d -cube, for which Theorem 5 gives the optimal bound and this bound can be extended to $k \in O(1)$. Theorem 1 gives an upper bound of $O(d^2k)$ on $h(\mathcal{T}(G,s))$, which is optimal for $d \in O(1)$. The rest of this section is devoted to proving the following result on the first-passage-percolation cover time of the (d,k) -grid, which gives an optimal bound on the height of $\mathcal{T}(G,s)$ for all values of k and d .

Theorem 6. *Let G be the (d,k) -grid and $n = (k+1)^d$. Then, for any vertex $s \in V(G)$, we have that $\tau(G,s) = O(k)$ with probability $1 - o_n(1)$.*

Before jumping into the proof, we note that applying Lemma 2 gives the following corollary of Theorem 6.

Corollary 2. *Let G be the (d,k) -grid and $n = (k+1)^d$. For any vertex $s \in V(G)$ we have that with probability $1 - o_n(1)$, $h(\mathcal{T}(G,s)) = \Theta(dk)$.*

To prove Theorem 6, we will make use of a concentration result about the first-passage percolation time on the d -cube.

Lemma 6. *Let Q be the d -cube, let $s \in V(Q)$ be arbitrary, and let $\bar{s} \in V(Q)$ be the unique vertex at distance d from s . Then there exist universal constants $c > 0$ and $x_0 > 0$ such that,*

$$\Pr\{\tau(Q, s, \bar{s}) > x\} \leq e^{-cx} ,$$

for all $x \geq x_0$.

Proof. We assume that d is greater than some sufficiently large constant, d_0 . Otherwise the result follows trivially from the union bound: With probability at least $1 - d2^{d-1}e^{-x/d}$, every edge of the d -cube has weight at most x/d . For $d \leq d_0$, this satisfies the statement of the lemma with $x_0 = 3d_0 \ln d_0$ and $c = 1/(3d_0^2)$.

We will prove the result for all $x \leq d^2$. Proving it for x in this range is sufficient, by a standard bootstrapping argument: For $x > d^2$, let $r = 2^{\lceil \log(x/d^2) \rceil}$, so that

$$x_0 \leq x/r \leq d^2$$

where the first inequality holds provided that $d^2 \geq 2x_0$. Consider a modified version of Richardson's infection model, which has the same rules as the original process except that, for each $i \in \mathbb{N}$, if the process has not infected \bar{s} by time ix/r , then we restart the process from the beginning. Clearly the time to infect \bar{s} in this modified process dominates the time to infect \bar{s} in the original process, so

$$\Pr\{\tau(Q, s, \bar{s}) > x\} \leq \Pr\{\tau(G, s, \bar{s}) > x/r\}^r \leq (e^{-cx/r})^r = e^{-cx} .$$

Thus, it suffices to prove the lemma for $x_0 \leq x \leq d^2$. For each $i \in \{0, 1, \dots, d\}$, let L_i denote the subset of $\binom{d}{i}$ vertices whose distance to s is i . Balister et al. [2, Lemma 4] show that there are constants $\alpha, \gamma > 0$ such that, if we sample each edge of Q independently with probability α/d then, with probability at least $1 - e^{-\gamma d^2}$, there is a path of length $d - 4$ consisting entirely of sampled edges and having one endpoint in L_2 and one endpoint in L_{d-2} .

In our setting, where edge weights are independent exponential(1), if we only consider edges of weight at most $\ln(d/(d - \alpha))$, then we obtain a sample in which each edge is independently sampled with probability α/d . Therefore, with probability at least $1 - e^{-\gamma d^2}$, there is a path of weight at most $d \ln(d/(d - \alpha)) = O(1)$ joining a vertex u in L_2 to a vertex w in L_{d-2} .

Now, there are d edge-disjoint paths of length 2 joining s to u . Consider one such path, P . If P has weight greater than $x/3$ then at least one of P 's edges has weight greater than $x/6$, which occurs with probability at most $2e^{-x/6}$. Therefore, the probability that all d paths have weight greater than $x/3$ is at most $(2e^{-x/6})^d \leq e^{-adx}$ for $0 < a < 1/6 - \ln 2/x$. Similarly, with probability at least $1 - e^{-adx}$, there is a path of length 2 and weight at most $x/3$ joining w to \bar{s} .

Therefore,

$$\Pr\{\tau(Q, s, \bar{s}) > 2x/3 + d \ln(d/(d - a))\} \leq \Pr\{\tau(G, s, \bar{s}) > x\} \leq 2e^{-adx} + e^{-\gamma d^2}$$

provided that $x \geq \max\{3d \ln(d/(d - a)), \ln 2/(1/6 - a)\}$. This holds, for example, when $x \geq x_0 = 12 \ln 2$, $a = 1/12$, and $d \geq 1$. \square

Lemma 6 extends from $\tau(Q, s, \bar{s})$ to $\tau(Q, s, v)$ for any vertex $v \in V(Q)$ using the union bound and the fact that v is at distance at least $d/2$ from at least one of s or \bar{s} . (This argument is also used by Balister et al. [2].)

Corollary 3. *Let Q be the d -cube, and let $s, v \in V(Q)$ be arbitrary. Then there exist universal constants $c > 0$ and $x_0 > 0$ such that,*

$$\Pr\{\tau(Q, s, v) > x\} \leq e^{-cx_d} ,$$

for all $x \geq x_0$.

Corollary 3 says that the tail of $\tau(Q, s, v)$ is dominated by an exponential(cd) random variable. The next lemma shows that sums of independent copies of $\tau(Q, s, v)$ also behave (roughly) like sums of independent exponential(cd) random variables.

Lemma 7. *Let Q be a d -cube, let $s_1, \dots, s_k \in V(Q)$ and $v_1, \dots, v_k \in V(Q)$ be arbitrary, and let Z_1, \dots, Z_k be i.i.d., random variables where Z_i is distributed like $\tau(Q, s_i, v_i)$. Then there exist universal constants $c > 0$ and $x_0 > 0$ such that*

$$\Pr\left\{\sum_{i=1}^k Z_i > ak\right\} \leq e^{(x_0+1)k-acdk/2} ,$$

for all $a > 0$.

Proof. If X is exponential(cd), then Corollary 3 says that

$$\Pr\{\tau(Q, s_i, v_i) > x\} \leq e^{-cx_d} = \Pr\{X > x\}$$

for all $x > x_0$. This implies that

$$\Pr\{\tau(Q, s_i, v_i) > x\} \leq \Pr\{X + x_0 > x\}$$

for all $x > 0$, i.e., $X + x_0$ stochastically dominates $\tau(Q, s_i, v_i)$. Therefore, if X_1, \dots, X_k are independent exponential(1), then

$$\Pr\left\{\sum_{i=1}^k Z_i > ak\right\} \leq \Pr\left\{\sum_{i=1}^k X_i > (a - x_0)k\right\} \leq e^{(x_0+1)k-acdk/2} ,$$

where the second inequality is an application of (2). □

We can now finish the proof of Theorem 6.

Proof of Theorem 6. The idea of this proof is that, for any vertex v , there is a path from s to v that visits at most k d -cubes. Therefore, there is a path from s to v whose length can be expressed as a sum like that considered in Lemma 7.

For each vertex $u = (u_1, \dots, u_d)$ with $u_i \in \{0, \dots, k-1\}$ for each $i \in \{1, \dots, d\}$, define the subgraph $Q_u = G[V_u]$ of G induced by the vertex set

$$V_u = \{(u_1 + x_1, \dots, u_d + x_d) : x_i \in \{0, 1\} \text{ for each } i \in \{1, \dots, d\}\} .$$

Each Q_u is a $(d, 1)$ -grid, i.e, a d -cube. For any vertex $v \in G$, there is a sequence $v_1, v_2, \dots, v_{k'}$ of vertices in G with $k' \leq k$ such that

-
1. $s \in Q_{v_1}$ and $v \in Q_{v_{k'}}$;
 2. for each $i \in \{1, \dots, k' - 1\}$, Q_{v_i} and $Q_{v_{i+1}}$ have at least one vertex in common.
 3. for each $i \in \{1, \dots, k'\}$ and each $j \in \{1, \dots, k'\} \setminus \{i - 1, i, i + 1\}$, Q_{v_i} and Q_{v_j} have no edges (or vertices) in common.

The sequence $v_1, \dots, v_{k'}$ can be found with a greedy algorithm: Define $v'_0 = s$. Now, if v'_{i-1} and v differ in r coordinates, then there is some vertex v_i such that Q_{v_i} contains v'_{i-1} as well as some vertex v'_i whose distance to v is r less than the distance from v'_{i-1} to v . It is straightforward to verify that the resulting sequence of vertices $v_1, \dots, v_{k'}$ satisfies the three properties described above.

For each $i \in \{1, \dots, k'\}$, let $s_i = v'_{i-1}$ and $x_i = v'_i$. Now, observe that

$$\tau(G, s, v) \leq \sum_{i=1}^{k'} \tau(Q_{v_i}, s_i, x_i) .$$

Point 3, above, ensures that the random variables $\tau(Q_{v_1}, s_1, x_1), \dots, \tau(Q_{v_{k'}}, s_{k'}, x_{k'})$ can be partitioned into two sets of size $\lfloor k'/2 \rfloor$ and $\lceil k'/2 \rceil$ where the variables within each set are independent.

By Lemma 7 we now have

$$\begin{aligned} \Pr\{\tau(G, s, v) > ak\} &\leq \Pr\left\{\sum_{i=1}^{\lceil k'/2 \rceil} \tau(Q_{v_{2i-1}}, s_{2i-1}, x_{2i-1}) > ak\right\} + \Pr\left\{\sum_{i=1}^{\lfloor k'/2 \rfloor} \tau(Q_{v_{2i}}, s_{2i}, x_{2i}) > ak\right\} \\ &\leq 2e^{(x_0+1)k-acdk/2} . \end{aligned}$$

Applying the union bound over all $(k+1)^d$ choices of v completes the proof:

$$\begin{aligned} \Pr\{\tau(G, s) > ak\} &\leq \sum_{v \in V(G)} \Pr\{\tau(G, s, v) > ak\} \\ &\leq 2(k+1)^d e^{(x_0+1)k-acdk/2} \\ &= e^{d \ln(2(k+1)) + (x_0+1)k-acdk/2} \\ &= o_n(1) , \end{aligned}$$

for

$$a > \frac{2(d \ln(2(k+1)) + (x_0+1)k)}{cdk} . \quad \square$$

8 Lower Bounds for General Graphs

Next, we describe a series of lower bound constructions that match the upper bounds obtained in Theorems 1–3. In particular, these constructions show that the dependence on Δ in the upper bounds in the previous sections can not be asymptotically reduced.

In this section we prove the following theorem.




Figure 1: The graph H .

Theorem 7. *There exists a positive constant c such that for any given positive integers $1 < \Delta, D$ satisfying $D \geq 16e^3 \ln \Delta$, there exists an n -vertex graph G with maximum degree $\leq \Delta$, diameter $\leq D$, and a vertex s satisfying $\Pr\{h(\mathcal{T}(G, s)) \geq c(\Delta \ln n + \Delta D)\} \geq 1 - o_n(1)$.*

The graph G is obtained by gluing together two graphs H and I . The graph H has large diameter and high connectivity. The graph I has low connectivity and small diameter. By gluing them we obtain a graph of low diameter (because of I) but for which the infection is more likely to spread via H (because of its high connectivity), and hence will have a large height. We begin by defining and studying H and I individually.

8.1 The Ladder Graph H

Let L, δ, a be positive integers. The graph H is shown in Figure 1. The vertices of H are partitioned into L groups V_1, \dots, V_L , each of size δ . The edge set of H is

$$E(H) = \bigcup_{i=1}^{L-1} \{vw : v \in V_i, w \in V_{i+1}\} .$$

First we show that the infection spreads rather quickly in H , namely we prove upper bounds for $\tau(H, v, w)$.

Lemma 8. *Let $a > e^2$. Then for any $1 \leq i < j \leq n$ and any $v \in V_i, w \in V_j$ we have*

$$\Pr\{\tau(H, v, w) > 2aL/(e^2 \delta)\} \leq \exp(L - aL/(2e^2)) + \exp(-aL/(e^2 \delta)) .$$

Proof. Consider the following greedy algorithm for finding a path from v to w : The path starts at v (which is in V_i). When the path has reached some vertex $x \in V_k$, for $k < j - 1$, the algorithm extends the path by taking the minimum-weight edge joining x to some vertex in V_{k+1} . When the algorithm reaches some $x \in V_{j-1}$, it takes the edge xw .

Let $m = j - i$. Each of the first $m - 1$ edges in the resulting path has a weight that is the minimum of δ exponential(1) random variables. Thus, the sum of weights of these

edges is the sum of $m - 1$ exponential(δ) random variables, i.e. an Erlang($m - 1, \delta$) random variable. The weight of the final edge is an exponential(1) random variable. Thus we find

$$\begin{aligned} \Pr\{\tau(H, v, w) > 2aL/(e^2\delta)\} &\leq \Pr\{\text{Erlang}(m - 1, \delta) + \text{exponential}(1) > 2aL/(e^2\delta)\} \\ &\leq \Pr\{\text{Erlang}(m - 1, \delta) > aL/(e^2\delta)\} + \Pr\{\text{exponential}(1) > aL/(e^2\delta)\} \\ &\leq \Pr\{\text{Erlang}(L, \delta) > aL/(e^2\delta)\} + \exp(-aL/(e^2\delta)) \\ &\leq \exp(L - aL/(2e^2)) + \exp(-aL/(e^2\delta)). \end{aligned}$$

The first inequality follows from the discussion above. The second inequality follows from the union bound. The third inequality is because an Erlang(L, δ) random variable stochastically dominates an Erlang($m - 1, \delta$) random variable, and the definition of the exponential distribution. The final equality follows from the tail bound (2). \square

8.2 The Subdivided Tree I

Next, we consider a tree I that is obtained by starting with a perfect binary tree³ having L leaves and then subdividing each edge incident to a leaf $\lceil aL/\delta \rceil - 1$ times so that each leaf-incident edge becomes a path of length $\lceil aL/\delta \rceil$. Note that I has height $\lceil aL/\delta \rceil + \log_2 L - 1$ (we assume L is a power of 2).

We next show that the infection spreads rather slowly in I , namely we prove lower bounds for $\tau(I, v, w)$.

Lemma 9. *For any distinct leaves v and w we have $\Pr\{\tau(I, v, w) \leq 2aL/(e^2\delta)\} \leq \exp(-2aL/\delta)$.*

Proof. The path from v to w in I contains at least $2\lceil aL/\delta \rceil$ edges. Therefore, the weight of this path is lower-bounded by the sum of $2\lceil aL/\delta \rceil$ independent exponential(1) random variables. The lemma then follows by applying (1) to this sum. \square

8.3 Putting it Together

The lower-bound graph G is now constructed by taking a tree I with L leaves and a graph H with L groups V_1, \dots, V_L each of size $\delta = \lfloor (\Delta - 1)/2 \rfloor$. Next, we consider the leaves of I in the order they are encountered in a depth first-traversal of I and, for each $i \in \{1, \dots, L\}$ we identify the i th leaf of I with some vertex in V_i . See Figure 2.

Lemma 10. *For any vertex $s \in V_1$ in the graph G described above, we have*

$$\Pr\{h(\mathcal{T}(G, s)) < L - 1\} \leq L^2 \left(\exp((1 - a/2e^2)L) + \exp(-aL/(e^2\delta)) + \exp(-2aL/\delta) \right)$$

Proof. Recall that $\mathcal{T}(G, s)$ is the shortest-path tree rooted at s for the first-passage percolation in G . If this tree contains no edge of I , its height is at least $L - 1$. If it does use some

³A perfect binary tree, sometimes called a complete binary tree, is a binary tree in which all vertices have 0 or 2 children, and all leaves have the same depth: <https://xlinux.nist.gov/dads/HTML/perfectBinaryTree.html>




Figure 2: The lower bound graph G . Dashed segments denote subdivided edges (paths of length $\lceil aL/\delta \rceil$).

edge of I , then there must be two leaves v and w of I such that $\tau(I, v, w) \leq \tau(H, v, w)$. Since there are $\binom{L}{2} < L^2$ choices for the pair $\{v, w\}$, using Lemma 8 and Lemma 9, we can bound the probability of this event by

$$\begin{aligned} & L^2 (\Pr\{d_H(v, w) > 2aL/\delta\} + \Pr\{d_I(v, w) < 2aL/\delta\}) \\ & \leq L^2 \left(\exp((1 - a/2e^2)L) + \exp(-aL/(e^2\delta)) + \exp(-2aL/\delta) \right), \end{aligned}$$

which proves the lemma. \square

We now have all the ingredients to prove the main theorem of this section, Theorem 7.

Proof of Theorem 7. Let $a = 4e^2$, $\delta = (\Delta - 1)/2$, and let L be the largest power of 2 that is not larger than $D\Delta/8a$. Let G be the graph described above. The maximum degree of G is $2\delta + 1 = \Delta$, and the diameter of G is bounded by

$$2(aL/\delta + \log_2 L) \leq 2(a \times (D\Delta/8a)/(\Delta/2) + \log_2(D\Delta/8a)) \leq D,$$

and its number of vertices is

$$n = L\delta + (2L - 1) + L(aL/\delta - 1) < L(\delta + 1 + aL/\delta).$$

We have

$$L \geq D\Delta/4a = \Omega(D\Delta + \Delta \ln L + \Delta \ln(\delta + 1 + aL/\delta)) = \Omega(\Delta \ln n + \Delta D).$$

By Lemma 10, there exists a vertex s such that

$$\begin{aligned} \Pr\{h(\mathcal{T}(G, s) \geq \Omega(\Delta \ln n + \Delta D))\} &\geq \Pr\{h(\mathcal{T}(G, s) \geq L - 1)\} \\ &\geq 1 - L^2 \left(\exp((1 - a/2e^2)L) - \exp(-aL/(e^2\delta)) - \exp(-2aL/\delta) \right) \\ &= 1 - \left(\exp(-L + 2\ln L) - \exp(-8L/\Delta + 2\ln L) - \exp(-16e^2L/\Delta + 2\ln L) \right) = 1 - o_L(1) = 1 - o_n(1), \end{aligned}$$

completing the proof. \square

9 Lower Bounds for Degenerate Graphs

Theorem 7 shows that Theorem 1 cannot be strengthened without knowing more about G than its number of vertices, maximum degree, and diameter. Theorem 2 provides a stronger upper bound under the assumption that G is d -degenerate. In this section we show that Theorem 2 is also tight, even when restricted to very special subclasses of d -degenerate graphs.

First we show that the bound given by Theorem 2 for $O(1)$ -degenerate graphs is tight, even when we restrict our attention to planar graphs, which are 5-degenerate. Since planar graphs have genus 0, this lower bound also shows that Theorem 3, which applies to bounded genus graphs, is tight.

Theorem 8. *There exists an absolute constant $c > 0$ such that for any $\Delta > 1$ and $D \geq 10^6 \ln \Delta$ there exists a planar graph with diameter $\leq D$, maximum degree $\leq \Delta$, and a vertex s such that with probability $1 - o_n(1)$ we have $h(\mathcal{T}(G, s)) > c\sqrt{\Delta}(D + \ln n)$.*

Proof. Let $C = 10^5$, $a = e^2C$, $\delta = \Delta/2$, and $L = D\sqrt{\delta}/3a$, and Let H be the graph shown in Figure 3, where each V_i has δ vertices. Let I be the perfect binary tree with L leaves, with each leaf-incident edge subdivided $aL/\sqrt{\delta} - 1$ times. Let G be the graph obtained from identifying the i th leaf of I with an arbitrary vertex from V_i . Note that G is a planar graph with maximum degree $2\delta = \Delta$, diameter $2(aL/\sqrt{\delta} + 1 + \log_2 L) \leq D$, and $n = \delta L + L - 1 + (2L - 1) + L(aL/\delta - 1) = O(\delta L + L^2/\delta)$ vertices. Let s be an arbitrary vertex in V_1 . Since $L = \Omega(\sqrt{\Delta}(D + \ln n))$, to complete the proof, we need only show that with probability $1 - o_n(1)$ we have $h(\mathcal{T}(G, s)) \geq 2L - 2$.

Choose an arbitrary vertex $t \in V_L$. Let \mathcal{A} denote the event $\tau(H, s, t) \leq CL/\sqrt{\delta}$, and let \mathcal{B} denote the event “for all pairs v and w of leaves of I we have $\tau(I, v, w) > CL/\sqrt{\delta}$. Note that if both \mathcal{A} and \mathcal{B} happen, then the path in $\mathcal{T}(G, s)$ from s to t uses edges from H only, which implies the height of this tree is at least $2L - 2$. To complete the proof via the union bound, we need only show that each of \mathcal{A} and \mathcal{B} happen with probability $1 - o_L(1) = 1 - o_n(1)$.

We start with \mathcal{A} . In H , one can go from the vertex in-between V_i and V_{i+1} to the vertex in-between V_{i+1} and V_{i+2} by taking a path whose weight is distributed as a $Y_{\delta,1}$ random variable (recall the definition of a $Y_{a,b}$ random variable from Section 2). Therefore, we have

$$\tau(H, s, t) = X_1 + X_2 + \sum_{i=1}^{L-2} Z_i,$$




Figure 3: The graph H in the proof of Theorem 8.

where X_1, X_2 are independent exponential(1) random variables (weights of the first and last edges), and Z_i 's are independent $Y_{\delta,1}$ random variables. Since $C/3 \geq 3 \times (64 + 1024)$, Using Lemma 1 (concentration for the sum of $Y_{a,b}$ random variables) we have

$$1 - \Pr\{\mathcal{A}\} \leq 2\Pr\{X_1 > CL/3\sqrt{\delta}\} + \Pr\left\{\sum_{i=1}^{L-1} Z_i > CL/3\sqrt{\delta}\right\} \leq 2\exp(-CL/3\sqrt{\delta}) + \exp(-(L-2)/9) = o_L(1)$$

We now prove \mathcal{B} happens with high probability. The path connecting any pair of leaves of I contains at least $2aL/\sqrt{\delta}$ edges, each of them having an independent exponential(1) weight. Therefore, using union bound over all pairs and using (1) we get

$$1 - \Pr\{\mathcal{B}\} \leq \binom{L}{2} \times \Pr\{\text{Erlang}(2aL/\sqrt{\delta}, 1) \leq CL/\sqrt{\delta}\} \leq L^2 \times (eC/2a)^{2aL/\sqrt{\delta}} = o_L(1),$$

completing the proof. \square

Next we describe a lower-bound construction that is d -degenerate, has thickness d and treewidth $O(d)$. This construction shows that Theorem 2 is asymptotically tight for all values $d \leq \Delta$ (with certain restrictions etc.).

Theorem 9. *There exists an absolute constant $c > 0$ such that for any $\Delta > 1$ with $D \geq 10^6 \ln \Delta$ and $d < \Delta$, there exists a graph G with diameter $\leq D$, maximum degree $\leq \Delta$, and the following properties:*

- (i) G is d -degenerate, has thickness $\leq d$ and treewidth $\leq 2d + 1$.
- (ii) G has a vertex s such that with probability $1 - o_n(1)$ we have $h(\mathcal{T}(G, s)) > c\sqrt{d\Delta}(D + \ln n)$.

Proof. Let $C = 10^5$, $a = e^2C$, $\delta = \Delta/2$, and $L = D\sqrt{d\Delta}/8a$, and Let H be the graph shown in Figure 4, where each V_i has δ vertices and each V'_i has d vertices, and each of the pairs $(V_1, V'_1), (V'_1, V_2), (V_2, V'_2)$, etc. form a complete bipartite graph. Let I be the perfect binary tree with L leaves, with each leaf-incident edge subdivided $aL/\sqrt{d\delta} - 1$ times. Consider




Figure 4: The d -degenerate graph H used in the proof of Theorem 9. In this example, $\delta = 7$ and $d = 3$.

the leaves of I in the order they are encountered in a depth first-traversal, for each $i \in \{1, \dots, L\}$ identify the i th leaf of I with some vertex in V_i . Let G be the resulting graph. Note that G has maximum degree $2\delta = \Delta$, diameter $\leq 2(1 + aL/\sqrt{d\delta} + \log_2 L) \leq D$, and $n = (\delta + D)L + 2L - 1 + L(aL/d\delta - 1) = O(\Delta L + L^2/d\Delta)$ vertices.

(i) Graph G is d -degenerate because the vertices of degree greater than d form an independent set. Therefore, every induced subgraph of G is either an independent set (so has a vertex of degree 0) or contains a vertex of degree at most d .

To see that G has thickness d , for each $i = 1, \dots, L$, assign to each vertex of V'_i a distinct colour from one of d colour classes. Now partition the edges incident to these vertices among d subgraphs depending on the color of the vertex they are incident to. Edges not incident to these vertices can be assigned to any subgraph. With this partition of edges, each subgraph becomes a subgraph of the planar graph used in the proof of Theorem 8.

To see that G has treewidth $2d + 1$, we build a tree decomposition of G with bags of maximum size $2d + 2$. For convenience, we define $V_0 = V_{L+1} = \emptyset$.

We begin with a tree T' of empty bags that has the same shape as I . For each vertex v of I , let B_v denote the bag of v .

1. Assign each vertex of v of I to B_v and to the (up to 2) children of B_v in T' .
2. Let v_1, \dots, v_L be the leaves of I ordered so that each $v_i \in V_i$. In the leaf bag B_{v_i} of T' we add all vertices in V'_{i-1} and V'_i .

Now each vertex in V'_i appears in B_{v_i} and $B_{v_{i+1}}$; so we add all vertices of V'_i to each of the bags on the path in T' from B_{v_i} to $B_{v_{i+1}}$.

3. Finally, to each B_{v_i} we attach $\delta - 1$ bags as leaves of T' ; in each bag we put all the vertices in $V'_i \cup V'_{i+1}$, and a distinct vertex of $V_i \setminus \{v_i\}$. We call each such bag B_v , where v is the unique vertex of $V_i \setminus \{v_i\}$ contained in the bag.

No bag contains more than $2d + 2$ vertices: for a leaf v_i , B_{v_i} contains v_i and its parent, as well as vertices in $V'_{i-1} \cup V'_i$. For a non-leaf vertex v of I , observe that (in any

binary tree) there are at most two distinct indices i, j such that v lies on the (v_i, v_{i+1}) -path in I and on the (v_j, v_{j+1}) -path, hence B_v contains v and its parent, as well as possibly V'_i and V'_j . For each $v \in V_i \setminus \{v_i\}$, B_v contains at most $2d + 1$ vertices; v and the vertices in $V'_{i-1} \cup V'_i$.

For each edge vw of G , there is some bag that contains both v and w : If vw is an edge of T with v a child of w then B_v contains both v and w . Otherwise, $v \in V_i$ and $w \in V'_{i-1}$ or $w \in V'_i$, in which case v and w appear in B_v .

Finally, for each vertex v of G , the subgraph of T' induced by bags containing v is connected: For a vertex $v \in I$ this subgraph is either an edge or a single vertex. For a vertex $v \in V_i$ this subgraph is a single vertex. For a vertex $v \in V'_i$ this subgraph is a path joining two vertices of T' .

Therefore, T' is a tree-decomposition of G whose largest bag has size $2d + 2$, and thus treewidth of G is at most $2d + 1$.

(ii) Let s be an arbitrary vertex in V_1 . Since $L = \Omega(\sqrt{d}\Delta(D + \ln n))$, to prove part (ii) we need only show that with probability $1 - o_n(1)$ we have $h(\mathcal{T}(G, s)) \geq 2L - 2$.

Choose an arbitrary vertex $t \in V_L$. Let \mathcal{A} denote the event $\tau(H, s, t) \leq CL/\sqrt{d}\delta$, and let \mathcal{B} denote the event “for all pairs v and w of leaves of I we have $\tau(I, v, w) > CL/\sqrt{d}\delta$ ”. Note that if both \mathcal{A} and \mathcal{B} happen, then the path in $\mathcal{T}(G, s)$ from s to t uses edges from H only, which implies the height of this tree is at least $2L - 2$. To complete the proof via the union bound, we need only show that each of \mathcal{A} and \mathcal{B} happen with probability $1 - o_L(1) = 1 - o_n(1)$.

We start with \mathcal{A} . In H , one can go from a given vertex in V'_i to some vertex in V'_{i+1} by taking a path whose weight is distributed as a $Y_{\delta, d}$ random variable. Therefore, $\tau(H, s, t)$ is stochastically dominated by

$$X_1 + X_2 + \sum_{i=1}^{L-2} Z_i,$$

where X_1, X_2 are independent exponential(1) random variables (weights of the first and last edges), and Z_i 's are independent $Y_{\delta, d}$ random variables. Since $C/3 \geq 3 \times (64 + 1024)$, Using Lemma 1 (concentration for the sum of $Y_{a, b}$ random variables) we have

$$1 - \Pr\{\mathcal{A}\} \leq 2\Pr\{X_1 > CL/3\sqrt{d}\delta\} + \Pr\left\{\sum_{i=1}^{L-2} Z_i > CL/3\sqrt{d}\delta\right\} \leq 2\exp(-CL/3\sqrt{d}\delta) + \exp(-(L-2)/9) = o_L(1)$$

We now prove \mathcal{B} happens with high probability. The path connecting any pair of leaves of I contains at least $2aL/\sqrt{d}\delta$ edges, each of them having an independent exponential(1) weight. Therefore, using union bound over all pairs and using (1) we get

$$1 - \Pr\{\mathcal{B}\} \leq \binom{L}{2} \times \Pr\{\text{Erlang}(2aL/\sqrt{d}\delta, 1) \leq CL/\sqrt{d}\delta\} \leq L^2 \times (eC/2a)^{2aL/\sqrt{d}\delta} = o_L(1),$$

completing the proof. □

Acknowledgements

Some of this research took place at the Workshop on Random Geometric Graphs and Their Applications to Complex Networks, at the Banff International Research Station, November 6–11, 2016. More of this research took place at the Rio Workshop on Geometry and Graphs, at IMPA, February 12–18, 2017. In both cases, we are grateful to the workshop organizers and other participants for providing stimulating working environments. We are especially grateful to Gábor Lugosi for helpful discussions on many aspects of this work and to Tasos Sidiropoulos for asking us about planar graphs.

References

- [1] Antonio Auffinger, Michael Damron, and Jack Hanson. 50 years of first passage percolation. *arXiv*, 1511.03262 [math.PR], 2016.
- [2] P. N. Balister, B. Bollobás, A. M. Frieze, and O. M. Riordan. The first-passage diameter of the cube, 2017. Unpublished manuscript.
- [3] Béla Bollobás. The isoperimetric number of random regular graphs. *Eur. J. Comb.*, 9(3):241–244, 1988.
- [4] Béla Bollobás and Yoshiharu Kohayakawa. On Richardson’s model on the hypercube. In *Combinatorics, Geometry, and Probability (Cambridge 1993)*, pages 129–137. Cambridge University Press, 1997.
- [5] Stéphane Boucheron, Gábor Lugosi, and Pascal Massart. *Concentration inequalities: A nonasymptotic theory of independence*. Oxford university press, 2013.
- [6] Luc Devroye. Branching processes in the analysis of the heights of trees. *Acta Informatica*, 24(3):277–298, 1987.
- [7] R. Durrett. Stochastic growth models: recent results and open problems. In *Mathematical approaches to problems in resource management and epidemiology (Ithaca, NY, 1987)*, volume 81 of *Lecture Notes in Biomath.*, pages 308–312. Springer, Berlin, 1989.
- [8] James Allen Fill and Robin Pemantle. Percolation, first-passage percolation and covering times for Richardson’s model on the n -cube. *The Annals of Applied Probability*, 3(2):593–629, 1993.
- [9] Svante Janson. Tail bounds for sums of geometric and exponential variables. available in <http://www2.math.uu.se/~svante/papers/sjN14.pdf>.
- [10] Anders Martinsson. Unoriented first-passage percolation on the n -cube. *The Annals of Applied Probability*, 26(5):2597–2625, 2016.
- [11] Colin McDiarmid. Concentration. In Michel Habib, Colin McDiarmid, Jorge Ramirez-Alfonsin, and Bruce Reed, editors, *Probabilistic Methods for Algorithmic Discrete Mathematics*, pages 195–248. Springer Berlin Heidelberg, Berlin, Heidelberg,

1998. available at http://www.stats.ox.ac.uk/people/academic_staff/colin_mcdiarmid/?a=4139.

- [12] Bojan Mohar and Carsten Thomassen. *Graphs on Surfaces*. Johns Hopkins University Press, Baltimore, Maryland, 2001.
- [13] Boris Pittel. Note on the heights of random recursive trees and random m-ary search trees. *Random Struct. Algorithms*, 5(2):337–348, 1994.
- [14] Thomas Wolle, Arie M.C.A. Koster, and Hans L. Bodlaender. A note on contraction degeneracy. Technical Report UU-CS-2004-042, University of Utrecht, 2004.

A Proof of Lemma 1

We will use the following inequality, which holds for any positive integer k and any real number λ (see [9, Theorem 5.1(ii)]):

$$\Pr\{\text{Erlang}(k, 1) \geq \lambda k\} \leq \exp(1 - \lambda). \quad (4)$$

We will also use the following inequality, which holds for any binomial random variable X , and any $M \leq E[X]$ (see [11, Theorem 2.3(c)]):

$$\Pr\{X < M/2\} \leq \exp(-M/8). \quad (5)$$

We will use the following version of Bernstein’s inequality (see Theorem 2.10 and Corollary 2.11 in [5]).

Theorem 10 (Bernstein’s inequality). *Let X_1, \dots, X_m be non-negative independent random variables for which there exist v, c satisfying*

$$\sum_{i=1}^m E[X_i^p] \leq vp!c^{p-2}/2$$

for all positive integers $p \geq 2$. Then for any $t > 0$ we have

$$\Pr\left\{\sum_{i=1}^m (X_i - E[X_i]) \geq ct + \sqrt{2vt}\right\} \leq e^{-t},$$

and

$$\Pr\left\{\sum_{i=1}^m (X_i - E[X_i]) \geq t\right\} \leq \exp\left(-\frac{t^2}{2v + 2ct}\right),$$

We begin with a helper lemma.

Lemma 11. *For any t we have*

$$\Pr\{Y_{a,b} > t\} \leq \exp(-at/64) + \exp(-abt^2/1024).$$

Proof. First, consider the case $t > 4$. Note that there exist a independent root-to-leaf paths, the weight of each is Erlang(2, 1). Hence, using (4) and since $t \geq 4$,

$$\Pr\{Y_{a,b} > t\} \leq \Pr\{\text{Erlang}(2, 1) > t\}^a \leq (\exp(1-t/2))^a \leq (\exp(-t/4))^a = \exp(-at/4) \leq \exp(-at/64).$$

The case $t \leq 0$ is trivial, so we consider the case $0 \leq t \leq 4$. Note that for such t we have $1 - \exp(-t/2) \geq t/8$. We say a node in the tree *survives* if each of the edges on its path to the root have weight at most $t/2$. Note that $Y_{a,b} > t$ implies no node at level 2 survives. The probability that a node at level 1 (children of the root) survives is $1 - \exp(-t/2)$, so the number of surviving nodes at level 1, S_1 , is a binomial random variable with mean $a(1 - \exp(-t/2)) \geq at/8$. From (5) we have

$$\Pr\{S_1 < at/16\} \leq \Pr\{S_1 < E[S_1]/2\} \leq \exp(-E[S_1]/8) \leq \exp(-at/64).$$

Conditioned on $S_1 \geq at/16$, the number of surviving nodes at level 2, S_2 , is a binomial random variable with mean $S_1 b(1 - \exp(-t/2)) \geq abt^2/128$, so using (5) again we have

$$\begin{aligned} \Pr\{Y_{a,b} > t | S_1 \geq at/16\} &\leq \Pr\{S_2 = 0 | S_1 \geq at/16\} \leq \Pr\{S_2 < abt^2/256 | S_1 \geq at/16\} \\ &\leq \exp(-abt^2/1024), \end{aligned}$$

completing the proof. \square

We are now ready to prove Lemma 1. Let X_1, \dots, X_m be i.i.d. distributed as $Y_{a,b}$ for some a, b . Then we want to prove $E[X_1] = O(1/a + 1/\sqrt{ab})$ and moreover,

$$\Pr\left\{\sum_{i=1}^m X_i \geq 3m(64/a + 1024/\sqrt{ab})\right\} \leq \exp(-m/9).$$

Let $d_1 = a/64$ and $d_2 = ab/1024$. For any positive integer p , by Lemma 11 we have

$$E[X_1^p] = \int_0^\infty \Pr\{X_1 > t^{1/p}\} dt \leq \int_0^\infty \exp(-d_1 t^{1/p}) + \int_0^\infty \exp(-d_2 t^{2/p})$$

For any positive numbers c, α , we have

$$\int_0^\infty \exp(-ct^\alpha) dt = \int_0^\infty \exp(-x) \frac{x^{1/\alpha-1}}{\alpha c^{1/\alpha}} dx = \frac{c^{-1/\alpha}}{\alpha} \int_0^\infty e^{-x} x^{1/\alpha-1} = \frac{c^{-1/\alpha} \Gamma(1/\alpha)}{\alpha}, \quad (6)$$

whence,

$$E[X_1^p] \leq p d_1^{-p} \Gamma(p) + p d_2^{-p/2} \Gamma(p/2)/2$$

In particular, setting $p = 1$ gives $E[X_1] \leq 64/a + 1024/\sqrt{ab} =: c$. Let $v = 4c^2 m$. For $p \geq 2$, we have

$$\sum_{i=1}^m E[X_i^p] \leq m p d_1^{-p} \Gamma(p) + m p d_2^{-p/2} \Gamma(p/2)/2 \leq m p! d_1^{-p} + m p! d_2^{-p/2}/2 \leq v p! c^{p-2}/2.$$

Bernstein's inequality (Theorem 10) gives that for all t ,

$$\Pr\left\{\sum_{i=1}^m X_i \geq m(64/a + 1024/\sqrt{ab}) + ct + 3c\sqrt{mt}\right\} \leq e^{-t},$$

and choosing $t = m/9$ completes the proof of the lemma.