

Estimates and Computations for Melting and Solidification Problems

J. M. Greenberg[†]

Carnegie Mellon University
Department of Mathematical Sciences
Pittsburgh, PA 15213

1 Introduction

In this note we focus on rigorous estimates for melting and solidification problems. These estimates guarantee the convergence of solutions to nontrivial equilibrium patterns. We also present some interesting numerical simulations which demonstrate the equilibrium structures and the approach of the system to equilibrium. The novel feature of these calculations is the linking of the small parameter in the system, δ , to the grid spacing.

Our governing equations are an energy balance for the medium and an evolution equation for a nonconservative “order parameter,” p . In dimensionless variables the energy equation takes the form

$$T_t + (1 - p^2)^m p_t - a^2 \Delta T = 0, \quad (x, y) \in \Omega \quad (1.1)$$

while the evolution equation for p may be written as

$$\alpha \delta p_t - \lambda^2 \delta^2 \Delta p = \mu T (1 - p^2)^m + p (1 - p^2), \quad (x, y) \in \Omega. \quad (1.2)$$

Here Δ is the two-dimensional Laplacian and T is the dimensionless temperature. The parameters α, a, λ , and μ are positive and order 1, $0 < \delta \ll 1$, and m is a nonnegative integer. The equations with $m = 0$ are the most frequently studied, the case where $m = 2$ has been investigated by Sekerka and coworkers [1-3] and Almgren and Almgren [4]. For definiteness we take $m = 1$ and restrict our attention to the situation where Ω is the unit square

$$\Omega = \{(x, y) \mid -1/2 < x < 1/2 \text{ and } -1/2 < y < 1/2\}. \quad (1.3)$$

On $\partial\Omega$ we assume that the normal derivatives of T and p vanish. Initial conditions for T and p are prescribed. Noting that at a local maximum (minimum) of p that $\Delta p \leq 0$ ($\Delta p \geq 0$) it is easily checked that if the initial data for p satisfies $-1 \leq p(x, y, 0^+) \leq 1$, $(x, y) \in \bar{\Omega}$, then the same estimates obtain for all future $t > 0$. We shall exploit this fact in what follows.

The system (1.1) and (1.2) is customarily referred to as the diffuse-interface description of melting and solidification. There is a corresponding sharp-interface description which is obtained formally from (1.2) by letting δ tend to 0^+ . The derivation of this description may be found in Caginalp [5-6].

In the sharp-interface description p only takes on the values -1 and $+1$. Solid regions are those where $p = -1$ and liquid regions those where $p = +1$. In what follows

[†]This research was partially supported by the Applied Mathematical Sciences Program, U.S. Department of Energy and the Mathematics and Computer Science Division, Army Research Office.

$$\Gamma(t) = \{(x, y) \mid x = \hat{x}(s, t), y = \hat{y}(s, t), 0 \leq s \leq L\} \quad (1.4)$$

will be a curve separating liquid and solid regions and $\mathbf{n}(s, t)$ will be the unit normal to Γ which points into the liquid region. The unit tangent, \mathbf{t} , to Γ is chosen so that $\mathbf{n}(s, t) \times \mathbf{t}(s, t) = \mathbf{e}_3$. K is the curvature of Γ at $(\hat{x}, \hat{y})(s, t)$, and

$$c(s, t) \stackrel{def}{=} (\hat{x}_t, \hat{y}_t) \cdot \mathbf{n}(s, t) \quad (1.5)$$

is the speed at which Γ moves in the direction \mathbf{n} . A singular perturbation analysis of (1.1) and (1.2), with $m = 1$, yields the following results:

1. T is continuous across Γ ;
2. On Γ the Gibbs-Thomson relation

$$\alpha c + \lambda^2 \delta K = -2^{1/2} \lambda \mu T_M \quad (1.6)$$

holds; and finally across Γ

$$\frac{4}{3}c = a^2 \left(\frac{\partial T^-}{\partial n} - \frac{\partial T^+}{\partial n} \right). \quad (1.7)$$

In (1.6), $T_M(s) = T(\hat{x}(s, t), \hat{y}(s, t))$ is the local melt temperature while in (1.7), $\frac{\partial T^+}{\partial n}$ is the normal derivative of T in the liquid region and $\frac{\partial T^-}{\partial n}$ is the normal derivative of T in the solid region. In the sharp-interface description one solves the heat equation

$$T_t - a^2 \Delta T = 0 \quad (1.8)$$

in the portion of Ω away from interfaces. At equilibrium, $c = 0$ and (1.6) implies that $T_M = 0(\delta)$. In what follows, we shall restrict our attention to initial data $T(x, y, 0^+) = 0(\delta^{\frac{1}{2}})$.

The sharp-interface description becomes unduly complicated in the presence of multiple interfaces since all must be tracked. Difficulties also arise when interfaces merge. For these, and a plethora of other reasons, we focus on (1.1) and (1.2). Our principal results consist of a sequence of a-priori inequalities satisfied by solutions of the system (1.1) and (1.2). These estimates guarantee regularity of the solutions and that the solutions converge to equilibria as t tends to infinity.

One point worthy of note is it is possible to obtain solutions of (1.1) and (1.2) which approximate those generated by the sharp interface descriptions. These are produced by solving (1.1) and (1.2) with initial data $p^\delta(x, y, 0^+)$ which approximate the jump discontinuous data of the sharp interface theory but transition smoothly from -1 to $+1$ over an interval of width of $0(\delta)$ in the direction \mathbf{n} . For such

initial p 's (and initial temperature fields satisfying $T(x, y, 0^+) = 0(\delta^{\frac{1}{2}})$) we are guaranteed that terms like $\delta \int \int_{\Omega} |\nabla p^\delta|^2 dx dy$ and $\frac{1}{\delta} \int \int_{\Omega} (1 - p^2)^2 dx dy$ are bounded independently of δ and t and it is these estimates which link the solutions of the two theories.

We also note there is an “intermediate” model where one replaces (1.1) (here with $m = 1$) with

$$T(x, y, t) \equiv T_0(t) = h_0 - \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} (p - p^3/3)(x, y, t) dx dy \quad (1.9)$$

and solves

$$\alpha \delta p_t - \lambda^2 \delta^2 \Delta p = (\mu T_0 + p)(1 - p^2), \quad (x, y) \in \Omega \quad (1.10)$$

and

$$\frac{\partial p}{\partial n} = 0 \quad , \quad (x, y) \in \partial\Omega. \quad (1.11)$$

The constant h_0 in (1.9) is given by

$$h_0 = \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} (T + p - p^3/3)(x, y, 0^+) dx dy. \quad (1.12)$$

This “intermediate” model is justified by two facts. The first is that solutions of (1.1) and (1.2) satisfying the boundary conditions $\frac{\partial T}{\partial n} = \frac{\partial p}{\partial n} = 0$, $(x, y) \in \partial\Omega$ satisfy the overall energy balance:

$$h_0 \equiv \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} (T + p - p^3/3)(x, y, t) dx dy \quad (1.13)$$

for all $t \geq 0$ and the second is that the spatially varying portion of T decays to zero as t tends to infinity (for details see the comments following (2.17)).

2 Entropies and A-Priori Estimates

Our goal in this section is a set of inequalities of the form

$$\frac{\partial \eta}{\partial t} - \operatorname{div} \mathbf{q} \leq G \quad (2.1)$$

satisfied by solutions of

$$T_t + (1 - p^2) p_t - a^2 \Delta T = 0 \quad , \quad (x, y) \in \Omega, \quad (2.2)$$

$$\alpha\delta p_t - \lambda^2\delta^2\Delta p = (\mu T + p)(1 - p^2), \quad (x, y) \in \Omega, \quad (2.3)$$

and

$$\frac{\partial T}{\partial n} = \frac{\partial p}{\partial n} = 0, \quad (x, y) \in \partial\Omega. \quad (2.4)$$

Once again

$$\Omega = \{(x, y) \mid -1/2 < x < 1/2 \text{ and } -1/2 < y < 1/2\}. \quad (2.5)$$

For (2.1) to be of value we require

$$0 \leq \eta, \quad (x, y) \in \Omega \text{ and } \mathbf{q} \cdot \mathbf{n} = 0, \quad (x, y) \in \partial\Omega \quad (2.6)$$

and that either $G \leq 0$ in Ω or G is a-priori bounded in $L_1(\Omega \times (0, \infty))$. Throughout we exploit the pointwise estimates

$$-1 \leq p(x, y, t) \leq 1, \quad (x, y) \in \Omega \text{ and } t \geq 0 \quad (2.7)$$

satisfied by solutions of (2.3).

Once again we record the overall energy balance satisfied by solutions of (2.2) – (2.4), namely the identity:

$$\int_{-1/2}^{1/2} \int_{-1/2}^{1/2} (T + p - p^3/3)(x, y, t) dx dy \equiv h_0. \quad (1.13)$$

This is obtained by integrating (2.2) over Ω and exploiting (2.4)₁.

Our first inequality is obtained by multiplying (2.2) by $(\frac{\mu T}{\delta})$ and (2.3) by $\frac{p_t}{\delta}$ and adding the resulting identities. This yields (2.1) with

$$\eta_1 = \frac{\mu T^2}{2\delta} + \frac{\lambda^2\delta}{2} |\nabla p|^2 + \frac{(1 - p^2)^2}{4\delta}, \quad (2.8)$$

$$\mathbf{q}_1 = \frac{\mu a^2}{\delta} T \nabla T + \lambda^2\delta p_t \nabla p, \quad (2.9)$$

and

$$G_1 = -\alpha p_t^2 - \frac{\mu a^2 |\nabla T|^2}{\delta} \quad (2.10)$$

and (2.8)-(2.10) yield

$$\begin{aligned} & \frac{d}{dt} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} \left(\frac{\mu T^2}{2\delta} + \frac{\lambda^2 \delta}{2} |\nabla p|^2 + \frac{(1-p^2)^2}{4\delta} \right) (x, y, t) dx dy \\ & \leq - \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} \left(\alpha p_t^2 + \frac{\mu a^2}{\delta} |\nabla T|^2 \right) (x, y, t) dx dy. \end{aligned} \quad (2.11)$$

Once again we restrict our attention to initial data satisfying

$$\int_{-1/2}^{1/2} \int_{-1/2}^{1/2} \left(\frac{T^2}{2\delta} + \frac{\lambda^2 \delta}{2} |\nabla p|^2 + \frac{(1-p^2)^2}{4\delta} \right) (x, y, 0^+) dx dy = 0(1)$$

independent of δ .

If we next multiply (2.2) by αT_t we obtain

$$\frac{\alpha a^2}{2} \frac{\partial}{\partial t} |\nabla T|^2 - \alpha a^2 \operatorname{div} (T_t \nabla T) = -\alpha (T_t^2 + (1-p^2) p_t T_t)$$

and if we add this result to (2.1) with η_1 , \mathbf{q}_1 , G_1 we obtain

$$\frac{\partial \eta_2}{\partial t} - \operatorname{div} \mathbf{q}_2 \leq G_2$$

where

$$\eta_2 = \frac{\mu T^2}{2\delta} + \frac{\lambda^2 \delta}{2} |\nabla p|^2 + \frac{(1-p^2)^2}{4\delta} + \frac{\alpha a^2}{2} |\nabla T|^2 \quad (2.12)$$

$$\mathbf{q}_2 = \frac{\mu a^2}{\delta} T \nabla T + \lambda^2 \delta p_t \nabla p + \alpha a^2 T_t \nabla T \quad (2.13)$$

and

$$G_2 = -\alpha (p_t^2 + T_t^2 + (1-p^2) T_t p_t) - \frac{\mu a^2}{\delta} |\nabla T|^2 \leq \frac{-\alpha}{2} (p_t^2 + T_t^2) - \frac{\mu a^2}{\delta} |\nabla T|^2 \quad (2.14)$$

and (2.12)-(2.14) imply that

$$\begin{aligned} & \frac{d}{dt} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} \left(\frac{\mu T^2}{2\delta} + \frac{\lambda^2 \delta}{2} |\nabla p|^2 + \frac{(1-p^2)^2}{4\delta} + \frac{\alpha a^2}{2} |\nabla T|^2 \right) (x, y, t) dx dy \\ & \leq - \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} \left(\frac{\alpha}{2} (p_t^2 + T_t^2) + \frac{\mu a^2}{\delta} |\nabla T|^2 \right) (x, y, t) dx dy. \end{aligned} \quad (2.15)$$

If we now write

$$T(x, y, t) = T_0(t) + T_1(x, y, t) \quad , \quad (x, y) \in \Omega \quad (2.16)$$

where

$$T_0(t) = \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} T(x, y, t) dx dy \quad \text{and} \quad \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} T_1(x, y, t) dx dy \equiv 0, \quad (2.17)$$

then (2.15) implies that if T and p are in $H^1(\Omega)$ at $t = 0$, then they are in $H^1(\Omega)$ for all $t > 0$ and that T satisfies $\lim_{t \rightarrow \infty} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} T_1^2(x, y, t) dx dy = 0$.¹

If we make a similar decomposition of p , that is write

$$p(x, y, t) = p_0(t) + p_1(x, y, t) \quad (2.18)$$

where

$$p_0(t) = \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} p(x, y, t) dx dy \quad \text{and} \quad \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} p_1(x, y, t) dx dy = 0, \quad (2.19)$$

then (2.3) implies that

$$\alpha \delta \frac{dp_0}{dt} = \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} (\mu T + p)(1 - p^2)(x, y, t) dx dy \quad (2.20)$$

and

$$\alpha \delta \frac{d^2 p_0}{dt^2} = \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} (\mu (T_t (1 - p^2) - 2pT p_t) + (1 - 3p^2) p_t) (x, y, t) dx dy \quad (2.21)$$

¹Recall, $H^k(\Omega)$ consists of functions f satisfying

$$\|f\|_k^2 \stackrel{def}{=} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} \sum_{j=0}^k \left(\sum_{\substack{\alpha_1 + \alpha_2 = j \\ \alpha_i \geq 0}} \left(\frac{\partial^j f}{\partial x^{\alpha_1} \partial y^{\alpha_2}} \right)^2 \right) (x, y) dx dy < \infty$$

and $H^k(\Omega \times (0, s_*))$ consists of all functions f on $\Omega \times (0, s_*)$ satisfying

$$\|f\|_k^2 \stackrel{def}{=} \int_0^{s_*} \left(\int_{-1/2}^{1/2} \int_{-1/2}^{1/2} \sum_{j=0}^k \left(\sum_{\substack{\alpha_1 + \alpha_2 + \alpha_3 = j \\ \alpha_i \geq 0}} \left(\frac{\partial^j f}{\partial x^{\alpha_1} \partial y^{\alpha_2} \partial s^{\alpha_3}} \right)^2 \right) (x, y, s) dx dy \right) ds < \infty.$$

and (2.20) and (2.21) along with the fact that (2.15) implies that p_t and T_t are $L_2(\Omega \times (0, \infty))$ combine to yield the result that $\frac{dp_0}{dt}$ and $\frac{d^2p_0}{dt^2}$ are in $L_2(0, \infty)$ and that

$$\lim_{t \rightarrow \infty} \alpha \delta \frac{dp_0}{dt} = \lim_{t \rightarrow \infty} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} (\mu T + p)(1 - p^2)(x, y, t) dx dy = 0. \quad (2.22)$$

With the estimate (2.15) we can glean weak information about the long-term behavior of solutions of (2.2)-(2.4). In what follows we let

$$(T^t, p^t)(x, y, s) = (T, p)(x, y, t + s), \quad (x, y) \in \Omega \text{ and } 0 \leq s \leq s_*. \quad (2.23)$$

The inequality (2.15), together with (2.7), implies that for any s_* the functions T^t and p^t are in $H^1(\Omega \times (0, s_*))$ with bounds which depend only on the $H^1(\Omega)$ norms of $T(x, y, 0^+)$ and $p(x, y, 0^+)$ and s_* and that

$$\lim_{t \rightarrow \infty} \int_0^{s_*} \left(\int_{-1/2}^{1/2} \int_{-1/2}^{1/2} \left(\frac{\alpha}{2} \left((T_s^t)^2 + (p_s^t)^2 \right) + \frac{\mu a^2}{\delta} |\nabla T^t|^2 \right) (x, y, s) dx dy \right) ds = 0. \quad (2.24)$$

Moreover, (2.3) implies that $\lambda^2 \delta^2 \Delta p^t \in L_2(\Omega \times (0, s_*))$ with bounds which are independent of t . The boundedness of T^t and p^t in $H^1(\Omega \times (0, s_*))$ implies we can find an increasing sequence $\{t_i\}_{i=1}^{\infty}$ with $\lim_{i \rightarrow \infty} t_i = \infty$ and functions T^∞ and p^∞ in $H^1(\Omega \times (0, s_*))$ and Δp^∞ in $L_2(\Omega \times (0, s_*))$ with the following properties:

(T^∞, p^∞) is the strong $L_2(\Omega \times (0, s_*))$ limit of (T^{t_i}, p^{t_i}) while T_s^∞ , p_s^∞ , ∇T^∞ , ∇p^∞ , and Δp^∞ are the weak $L_2(\Omega \times (0, s_*))$ limits of the appropriate derivatives of T^{t_i} and p^{t_i} . Additionally, (2.24) implies

$$T_s^\infty = p_s^\infty = 0 \text{ and } \nabla T^\infty = 0 \text{ in } \Omega \times (0, s_*) \quad (2.25)$$

and thus that T^∞ is a constant and p^∞ is independent of s . The energy balance (1.13) yields

$$T^\infty = h_0 - \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} (p^\infty - (p^\infty)^3 / 3) (x, y) dx dy \quad (2.26)$$

while (2.7) implies that $-1 \leq p^\infty \leq 1$. (2.3) and (2.24) also imply that p^∞ satisfies the equilibrium equation

$$\lambda^2 \delta^2 \Delta p^\infty + (\mu T^\infty + p^\infty) (1 - (p^\infty)^2) = 0, \quad (x, y) \in \Omega \quad (2.27)$$

and

$$\frac{\partial p^\infty}{\partial n} = 0, \quad (x, y) \in \partial \Omega \quad (2.28)$$

where T^∞ is given by (2.26) and finally (2.27) and (2.28) imply that p^∞ is a critical point of the functional

$$N(p^\infty) \stackrel{def}{=} \frac{\mu(T^\infty)^2}{2\delta} + \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} \left(\frac{\lambda^2\delta}{2} |\nabla p^\infty|^2 + \frac{(1 - (p^\infty)^2)^2}{4\delta} \right) (x, y) dx dy. \quad (2.29)$$

Additionally, the functional

$$\eta(t, s_*) \stackrel{def}{=} \frac{1}{s_*} \int_0^{s_*} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} \left(\frac{\mu(T^t)^2}{2\delta} + \frac{\lambda^2\delta}{2} |\nabla p^t|^2 + \frac{(1 - (p^t)^2)^2}{4\delta} + \frac{\alpha a^2}{2} |\nabla T^t|^2 \right) (x, y, s) dx dy ds \quad (2.30)$$

is decreasing in t and has a limit η_∞ satisfying

$$N(p^\infty) \leq \eta_\infty. \quad (2.31)$$

To obtain stronger information we examine the time differentiated versions of (2.2)-(2.4), namely the equations:

$$T_{tt} + (1 - p^2)p_{tt} - 2pp_t^2 - a^2\Delta T_t - 0, \quad (x, y) \in \Omega, \quad (2.32)$$

$$\alpha\delta p_{tt} - \lambda^2\delta^2\Delta p_t = (\mu T_t + p_t)(1 - p^2) - 2p(\mu T + p)p_t, \quad (x, y) \in \Omega, \quad (2.33)$$

$$\frac{\partial T_t}{\partial n} = \frac{\partial p_t}{\partial n} = 0, \quad (x, y) \in \partial\Omega. \quad (2.34)$$

We shall also exploit

Lemma 1. Suppose $f \in H^1(\Omega)$. Then, for every $\epsilon > 0$

$$\int_{-1/2}^{1/2} \int_{-1/2}^{1/2} f^4(x, y) dx dy \leq 4 \left[\frac{(1 + \epsilon)}{\epsilon} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} f^2(x, y) dx dy + \epsilon \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} |\nabla f|^2(x, y) dx dy \right]^2.$$

We multiply (2.33) by p_t and make use of (2.7) to obtain

$$\frac{\alpha\delta}{2} \frac{\partial}{\partial t} p_t^2 - \lambda^2\delta^2 \operatorname{div} (p_t \nabla p_t) \leq -\lambda^2\delta^2 |\nabla p_t|^2 + \left(\frac{\mu T_t^2}{2} + \left(1 + \frac{\mu}{2}\right) p_t^2 \right) + 2\mu|T|p_t^2. \quad (2.35)$$

We note that (2.15) implies the term $\frac{\mu T_t^2}{2} + \left(1 + \frac{\mu}{2}\right) p_t^2$ is in $L_1(\Omega \times (0, \infty))$ and thus we confine our attention to $2\mu|T|p_t^2$. Schwarz's inequality and Lemma 1 imply

$$\begin{aligned} 2\mu \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} (|T|p_t^2)(x, y, t) dx dy &\leq 2\mu \left(\int_{-1/2}^{1/2} \int_{-1/2}^{1/2} T^2(x, y, t) dx dy \right)^{1/2} \left(\int_{-1/2}^{1/2} \int_{-1/2}^{1/2} p_t^4(x, y, t) dx dy \right)^{1/2} \\ &\leq 2\mu \left(\int_{-1/2}^{1/2} \int_{-1/2}^{1/2} T^2(x, y, t) dx dy \right)^{1/2} \left(\left(\frac{1+\epsilon}{\epsilon} \right) \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} p_t^2(x, y, t) dx dy + \epsilon \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} |\nabla p_t|^2(x, y, t) dx dy \right). \end{aligned}$$

Moreover, (2.11) guarantees that

$$\left(\int_{-1/2}^{1/2} \int_{-1/2}^{1/2} T^2(x, y, t) dx dy \right)^{1/2} \leq \left(\frac{2\delta}{\mu} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} \eta_1(x, y, 0^+) dx dy \right)^{1/2} \quad \begin{matrix} (2.36) \\ (2.37) \end{matrix}$$

and this guarantees that we may choose ϵ so that

$$2\mu\epsilon \left(\int_{-1/2}^{1/2} \int_{-1/2}^{1/2} T^2(x, y, t) dx dy \right)^{1/2} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} |\nabla p_t|^2(x, y, t) \leq \frac{\lambda^2 \delta^2}{2} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} |\nabla p_t|^2(x, y, t) dx dy. \quad (2.38)$$

With ϵ so chosen, (2.35) implies that

$$\begin{aligned} \frac{\alpha\delta}{2} \frac{d}{dt} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} p_t^2(x, y, t) &\leq \frac{-\lambda^2 \delta^2}{2} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} |\nabla p_t|^2(x, y, t) dx dy + \frac{\mu}{2} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} T_t^2(x, y, t) dx dy \\ &+ \left(1 + \frac{\mu}{2} + 2^{3/2} \left(\mu\delta \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} \eta_1(x, y, 0^+) dx dy \right)^{1/2} \left(1 + \frac{1}{\epsilon} \right) \right) \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} p_t^2(x, y, t) dx dy \end{aligned} \quad (2.39)$$

and (2.39) then implies that if

$$p_t(x, y, 0^+) = \frac{1}{\alpha\delta} (\lambda^2 \delta^2 \Delta p + (\mu T + p)(1 - p^2))(x, y, 0^+)$$

is in $L_2(\Omega)$, then p_t and thus Δp are in $L_2(\Omega)$ independently of t and $\nabla p_t \in L_2(\Omega \times (0, \infty))$.

The preceding estimate points up the advantage of the "intermediate" model discussed briefly in (1.9)-(1.12) over the full system. For the intermediate model the contankerous term $2\mu|T|p_t^2$ in (2.35) would have been replaced by $2\mu|T_0|p_t^2$ and this latter term is easily controlled since we know by our previous estimates that

$$|T_0| \leq \left(\frac{2\delta}{\mu} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} \eta_1(x, y, 0^+) dx dy \right)^{1/2}$$

and thus we would not have to invoke Lemma 1 and choose $\epsilon = 0$ ($\delta^{3/2}$). This latter choice, pumped up the coefficient of $\int_{-1/2}^{1/2} \int_{-1/2}^{1/2} p_t^2(x, y, t) dx dy$ in (2.39) to be 0 ($1/\delta^{3/2}$) which is far from optimal. If we chose to add some multiple of (2.39) to (2.15) and wanted the right hand side of the resulting inequality to be negative definite our multiplier would have to be $0(\delta^{3/2})$, an unacceptably small number. For the “intermediate” model we could use an $0(1)$ multiplier (for details see (2.61)-(2.63)).

On the other hand, if we multiply (2.32) by $(T_t + (1 - p^2)p_t)/a^2$ we find that

$$\begin{aligned} \frac{1}{2a^2} \frac{\partial}{\partial t} (T_t + (1 - p^2)p_t)^2 &= \frac{a^2}{2} \frac{\partial}{\partial t} (\Delta T)^2 \\ &= \operatorname{div} ((T_t + (1 - p^2)p_t) \nabla T_t) - |\nabla T_t|^2 - (1 - p^2) \nabla T_t \cdot \nabla p_t + 2pp_t \nabla T_t \cdot \nabla p \end{aligned} \quad (2.40)$$

and (2.40) implies that

$$\begin{aligned} \frac{a^2}{2} \frac{d}{dt} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} (\Delta T)^2(x, y, t) dx dy &\leq \\ -\frac{1}{2} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} |\nabla T_t|^2(x, y, t) dx dy &+ \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} |\nabla p_t|^2(x, y, t) dx dy \\ +4 \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} p_t^2 |\nabla p|^2(x, y, t) dx dy. & \end{aligned} \quad (2.41)$$

The first term on the right hand side of (2.41) is in controllable by virtue of the minus sign and the second term is $L_1(0, \infty)$ by virtue of (2.39). Thus we can confine our attention to the third term. To show this term is $L_1(0, \infty)$ we note that

$$\int_{-1/2}^{1/2} \int_{-1/2}^{1/2} p_t^2 |\nabla p|^2(x, y, t) dx dy \leq \left(\int_{-1/2}^{1/2} \int_{-1/2}^{1/2} p_t^4(x, y, t) dx dy \right)^{1/2} \left(\int_{-1/2}^{1/2} \int_{-1/2}^{1/2} |\nabla p|^4(x, y, t) dx dy \right)^{1/2}. \quad (2.42)$$

Lemma 1 then guarantees that

$$\left(\int_{-1/2}^{1/2} \int_{-1/2}^{1/2} p_t^4(x, y, t) dx dy \right)^{1/2} \leq \frac{(1 + \sqrt{5})}{2} \left(\int_{-1/2}^{1/2} \int_{-1/2}^{1/2} (p_t^2 + |\nabla p_t|^2)(x, y, t) dx dy \right) \quad (2.43)$$

and that

$$\begin{aligned}
& \left(\int_{-1/2}^{1/2} \int_{-1/2}^{1/2} |\nabla p|^4(x, y, t) dx dy \right)^{1/2} \leq \left(\int_{-1/2}^{1/2} \int_{-1/2}^{1/2} p_x^4(x, y, t) dx dy \right)^{1/2} + \left(\int_{-1/2}^{1/2} \int_{-1/2}^{1/2} p_y^4(x, y, t) dx dy \right)^{1/2} \\
& \leq \frac{(1 + \sqrt{5})}{2} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} (p_x^2 + p_y^2 + p_{xx}^2 + p_{yy}^2 + 2p_{xy}^2)(x, y, t) dx dy \\
& \leq \frac{(1 + \sqrt{5})}{2} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} (p_x^2 + p_y^2 + 4(\Delta p)^2)(x, y, t) dx dy
\end{aligned} \tag{2.44}$$

and (2.43) and (2.44) together with (2.41) guarantee that if $T(x, y, 0^+)$ and $p(x, y, 0^+)$ are in $H^2(\Omega)$ then $T(x, y, t)$ is in $H^2(\Omega)$ for all future times and additionally that $\nabla T_t \in L_2(\Omega \times (0, \infty))$.

The estimates (2.39) and (2.44) also guarantee that if $T(x, y, 0^+)$ and $p(x, y, 0^+)$ are in $H^2(\Omega)$, then the resulting orbit is compact in $H^1(\Omega)$ and that the w -limit set is nonempty in $H^1(\Omega)$. The fact that $\nabla T_t \in L_2(\Omega \times (0, \infty))$ further guarantees that the function T_1 defined in (2.16) and (2.17) converges to zero strongly in $H^1(\Omega)$.

Our final estimates pertain to solutions generated by initial data $T(x, y, 0^+) \in H^2(\Omega)$ and $p(x, y, 0^+) \in H^3(\Omega)$. These are obtained by multiplying (2.33) by p_{tt} . The resulting inequality is

$$\begin{aligned}
& \frac{\lambda^2 \delta^2}{2} \frac{\partial}{\partial t} |\nabla p_t|^2 + \alpha \delta p_{tt}^2 - \lambda^2 \delta^2 \operatorname{div} (p_{tt} \nabla p_t) = \\
& \mu (T_t(1 - p^2) - 2pT p_t) p_{tt} + (1 - 3p^2) p_t p_{tt} \\
& \leq \frac{\alpha \delta}{2} p_{tt}^2 + \frac{2\mu^2}{\alpha \delta} (T_t^2 + 4T^2 p_t^2) + \frac{8}{\alpha \delta} p_t^2.
\end{aligned} \tag{2.45}$$

Our preceding calculations imply that if $T(x, y, 0^+) \in H^2(\Omega)$ and $p(x, y, 0^+) \in H^3(\Omega)$, then

$$\frac{2\mu^2}{\alpha \delta} (T_t^2 + 4T^2 p_t^2) + \frac{8}{\alpha \delta} p_t^2$$

is in $L_1(\Omega \times (0, \infty))$ and thus (2.45) implies that $\nabla p_t \in L_2(\Omega)$ independently of t and that $p_{tt} \in L_2(\Omega \times (0, \infty))$. The former conclusion, together with (2.3), implies that $p \in H^3(\Omega)$ with bounds which are independent of t while the latter conclusion implies that

$$\lim_{t \rightarrow \infty} \alpha \delta p_t = \lim_{t \rightarrow \infty} (\lambda^2 \delta^2 \Delta p + (\mu T + p)(1 - p^2)) = 0$$

and that this limit is obtained strongly in $L_2(\Omega)$.

We conclude this section with some observations about the ‘‘intermediate’’ model introduced in (1.9)-(1.12). Once again in that model one replaces (2.3) with

$$\alpha \delta p_t - \lambda^2 \delta^2 \Delta p = (\mu T_0 + p)(1 - p^2), (x, y) \in \Omega \tag{2.46}$$

and

$$\frac{\partial p}{\partial n} = 0 \quad , \quad (x, y) \in \partial\Omega \quad (2.47)$$

where

$$T_0(t) = h_0 - \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} (p - p^3/3) (x, y, t) dx dy \quad \text{and} \quad h_0 = \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} (T + p - p^3/3) (x, y, 0^+) dx dy. \quad (2.48)$$

Solutions of this latter system also satisfy the pointwise estimates $-1 \leq p(x, y, t) \leq 1$ and a set of “entropy” inequalities similar to those satisfied by solutions of the full system. The most basic of these is

$$\frac{\partial \tilde{\eta}_1}{\partial t} - \operatorname{div} \tilde{\mathbf{q}}_1 = \tilde{G}_1 \quad (2.49)$$

where

$$\tilde{\eta}_1 = \frac{\mu T_0^2}{2\delta} + \frac{\lambda^2 \delta}{2} |\nabla p|^2 + \frac{(1 - p^2)^2}{4\delta} \quad (2.50)$$

$$\tilde{\mathbf{q}}_1 = \lambda^2 \delta p_t \nabla p \quad (2.51)$$

and

$$\tilde{G}_1 = -\alpha p_t^2. \quad (2.52)$$

Moreover, (2.49) - (2.52) imply that

$$\begin{aligned} & \frac{d}{dt} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} \left(\frac{\mu T_0^2}{2\delta} + \frac{\lambda^2 \delta}{2} |\nabla p|^2 + \frac{(1 - p^2)^2}{4\delta} \right) (x, y, t) dx dy \\ & \leq -\alpha \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} p_t^2(x, y, t) dx dy. \end{aligned} \quad (2.53)$$

The time differentiated identities

$$\alpha \delta p_{tt} - \lambda^2 \delta^2 \Delta p_t = \mu \left(\dot{T}_0 (1 - p^2) - 2p T_0 p_t \right) + (1 - 3p^2) p_t, \quad (2.54)$$

$$\dot{T}_0 = - \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} (1 - p^2) p_t(x, y, t) dx dy, \quad (2.55)$$

and

$$\frac{\partial p_t}{\partial n} = 0 \quad , \quad (x, y) \in \partial\Omega. \quad (2.56)$$

also yields:

$$\begin{aligned} & \frac{\alpha\delta}{2} \frac{\partial p_t^2}{\partial t} - \lambda^2 \delta^2 \operatorname{div} (p_t \nabla p_t) = \\ & -\lambda^2 \delta^2 |\nabla p_t|^2 + \mu \left(\dot{T}_0 (1 - p^2) - 2pT_0 p_t \right) p_t + (1 - 3p^2) p_t^2 \\ & \leq -\lambda^2 \delta^2 |\nabla p_t|^2 + \frac{\mu}{2} \left(\int_{-1/2}^{1/2} \int_{-1/2}^{1/2} p_t^2(x, y, t) dx dy \right) \\ & + 2^{3/2} \mu^{1/2} \delta^{1/2} \left(\int_{-1/2}^{1/2} \int_{-1/2}^{1/2} \tilde{\eta}_1(x, y, 0^+) dx dy \right)^{1/2} p_t^2 + \left(1 + \frac{\mu}{2}\right) p_t^2 \end{aligned} \quad (2.57)$$

and

$$\begin{aligned} & \frac{\lambda^2 \delta^2}{2} \frac{\partial |\nabla p_t|^2}{\partial t} - \lambda^2 \delta^2 \operatorname{div} (p_{tt} \nabla p_t) = \\ & -\alpha\delta p_{tt}^2 + \mu \left(\dot{T}_0 (1 - p^2) - 2pT_0 p_t \right) p_{tt} + (1 - 3p^2) p_t p_{tt} \\ & \leq -\frac{\alpha\delta}{2} p_{tt}^2 + \frac{1}{\alpha\delta} \left(2\mu^2 \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} p_t^2(x, y, t) dx dy + \left(4 + 16\mu\delta \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} \tilde{\eta}_1(x, y, 0^+) dx dy \right) p_t^2 \right). \end{aligned} \quad (2.58)$$

The former follows from multiplying (2.54) by p_t and the latter from multiplying (2.54) by p_{tt} . Moreover, if we let

$$\beta_2 = \left(1 + \mu + 2^{3/2} \mu^{1/2} \delta^{1/2} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} \tilde{\eta}_1(x, y, 0^+) dx dy \right), \quad (2.59)$$

$$\beta_3 = 4 + 2\mu^2 + 16\mu\delta \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} \tilde{\eta}_1(x, y, 0^+) dx dy, \quad (2.60)$$

$$\tilde{\eta}_2 = \tilde{\eta}_1 + \frac{\alpha^2 \delta}{8\beta_2} p_t^2 = \frac{\mu T_0^2}{2\delta} + \frac{\lambda^2 \delta}{2} |\nabla p|^2 + \frac{(1-p^2)^2}{4\delta} + \frac{\alpha^2 \delta}{8\beta_2} p_t^2, \quad (2.61)$$

$$\tilde{\mathbf{q}}_2 = \tilde{\mathbf{q}}_1 + \frac{\alpha \lambda^2 \delta^2}{4\beta_2} p_t \nabla p_t = \lambda^2 \delta p_t \nabla p + \frac{\alpha \lambda^2 \delta^2}{4\beta_2} p_t \nabla p_t, \quad (2.62)$$

$$\begin{aligned} \tilde{G}_2 &= \tilde{G}_1 + \frac{\alpha}{4\beta_2} \left(1 + \frac{\mu}{2} + 2^{3/2} \mu^{1/2} \delta^{1/2} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} \tilde{\eta}_1(x, y, 0^+) dx dy \right) p_t^2 \\ &+ \frac{\alpha \mu}{8\beta_2} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} p_t^2(x, y, t) dx dy - \frac{\alpha \lambda^2 \delta^2}{4\beta_2} |\nabla p_t|^2, \end{aligned} \quad (2.63)$$

$$\tilde{\eta}_3 = \tilde{\eta}_2 + \frac{\alpha^2 \lambda^2 \delta^3}{8\beta_3} |\nabla p_t|^2 = \frac{\mu T_0^2}{2\delta} + \frac{\lambda^2 \delta}{2} |\nabla p|^2 + \frac{(1-p^2)^2}{4\delta} + \frac{\alpha^2 \delta}{8\beta_2} p_t^2 + \frac{\alpha^2 \lambda^2 \delta^3}{8\beta_3} |\nabla p_t|^2 \quad (2.64)$$

$$\tilde{\mathbf{q}}_3 = \tilde{\mathbf{q}}_2 + \frac{\alpha^2 \lambda^2 \delta^3}{4\beta_3} p_{tt} \nabla p_t = \lambda^2 \delta p_t \nabla p + \frac{\alpha \lambda^2 \delta^2}{4\beta_2} p_t \nabla p_t + \frac{\alpha^2 \lambda^2 \delta^3}{4\beta_3} p_{tt} \nabla p_t, \quad (2.65)$$

and

$$\begin{aligned} \tilde{G}_3 &= \tilde{G}_2 + \frac{\alpha}{4\beta_3} \left(4 + 16\mu\delta \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} \tilde{\eta}_1(x, y, 0^+) dx dy \right) p_t^2 \\ &+ \frac{\alpha}{2\beta_3} \mu^2 \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} p_t^2(x, y, t) dx dy - \frac{\alpha^2 \delta^2}{8\beta_3} p_{tt}^2 \end{aligned} \quad (2.66)$$

we obtain the inequalities

$$\frac{\partial \tilde{\eta}_2}{\partial t} - \operatorname{div} \tilde{\mathbf{q}}_2 \leq \tilde{G}_2 \quad \text{and} \quad \frac{\partial \tilde{\eta}_3}{\partial t} = \operatorname{div} \tilde{\mathbf{q}}_3 \leq \tilde{G}_3 \quad (2.67)$$

and these imply

$$\begin{aligned} &\frac{d}{dt} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} \left(\frac{\mu T_0^2}{2\delta} + \frac{\lambda^2 \delta}{2} |\nabla p|^2 + \frac{(1-p^2)^2}{4\delta} + \frac{\alpha^2 \delta}{8\beta_2} p_t^2 \right) (x, y, t) dx dy \\ &\leq -\frac{3\alpha}{4} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} p_t^2(x, y, t) dx dy - \frac{\alpha \lambda^2 \delta^2}{4\beta_2} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} |\nabla p_t|^2(x, y, t) dx dy, \end{aligned} \quad (2.68)$$

and

$$\begin{aligned}
& \frac{d}{dt} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} \left(\frac{\mu T_0^2}{2\delta} + \frac{\lambda^2 \delta |\nabla p|^2}{2} + \frac{(1-p^2)^2}{4\delta} + \frac{\alpha \delta}{8\beta_2} p_t^2 + \frac{\alpha^2 \lambda^2 \delta^3}{8\beta_3} |\nabla p_t|^2 \right) (x, y, t) dx dy \\
& \leq -\frac{\alpha}{2} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} p_{tt}^2(x, y, t) - \frac{\alpha \lambda^2 \delta^2}{4\beta_2} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} |\nabla p_t|^2(x, y, t) dx dy \\
& \quad - \frac{\alpha^2 \delta^2}{4\beta_3} \int_{-1/2}^{1/2} \int_{-1/2}^{1/2} p_{tt}^2(x, y, t) dx dy.
\end{aligned} \tag{2.69}$$

We leave it to the reader to supply the precise conclusions derivable from the inequalities (2.53), (2.63), and (2.69). The key point of these inequalities is that the parameter δ appears optimally.

3 Computational Experiments

In this section we present some numerical simulations of the system (2.2) - (2.4). We let N be an integer and set

$$dx = dy = 1/N. \tag{3.1}$$

Our approximate solutions will be piecewise constant on the cells

$$\Omega_{i,j} = \left\{ (x, y) \mid -\frac{1}{2} + \frac{(i-1)}{N} < x < -\frac{1}{2} + \frac{i}{N} \text{ and } -\frac{1}{2} + \frac{(j-1)}{N} < y < -\frac{1}{2} + \frac{j}{N} \right\} \tag{3.2}$$

where $1 \leq i, j \leq N$ and thus the cell averages $(T_{i,j}, p_{i,j})$ may also be thought of as the point values at the cell centers

$$(x_i, y_j) = \left(-\frac{1}{2} + \frac{2i-1}{2N}, -\frac{1}{2} + \frac{2j-1}{2N} \right). \tag{3.3}$$

We impose the boundary conditions

$$\frac{\partial T}{\partial n} = \frac{\partial p}{\partial n} = 0 \quad , \quad (x, y) \in \partial\Omega \tag{3.4}$$

by insisting that the extended grid functions satisfy

$$\left. \begin{aligned}
(T_{0,k}, p_{0,k}) &= (T_{1,k}, p_{1,k}), \\
(T_{N+1,k}, p_{N+1,k}) &= (T_{N,k}, p_{N,k}), \\
(T_{k,0}, p_{k,0}) &= (T_{k,1}, p_{k,1}), \\
\text{and} \\
(T_{k,N+1}, p_{k,N+1}) &= (T_{k,N}, p_{k,N})
\end{aligned} \right\} \quad (3.5)$$

for $1 \leq k \leq N$.

For any function u defined on the grid and satisfying (3.5) we let

$$\Delta_N u_{i,j} = N^2 (u_{i+1,j} + u_{i-1,j} + u_{i,j+1} + u_{i,j-1} - 4u_{i,j}) \quad (3.6)$$

be the discrete laplacian applied to u . For functions u and v satisfying (3.5) we define

$$[u, v] = \sum_{i,j=1}^N u_{i,j} v_{i,j} / N^2 \quad (3.7)$$

and note that

$$-[u, \Delta_N u] = \sum_{i=1}^N \sum_{j=1}^{N-1} (u_{i,j+1} - u_{i,j})^2 + \sum_{j=1}^N \sum_{i=1}^{N-1} (u_{i+1,j} - u_{i,j})^2. \quad (3.8)$$

Such grid functions also satisfy the discrete version of Lemma 1, namely the inequality

$$[u^2, u^2] = [u^4, 1] \leq 4 \left(\frac{(1+\epsilon)}{\epsilon} [u, u] - \epsilon [u, \Delta_N u] \right)^2 \quad (3.9)$$

for every $\epsilon > 0$.

Finally, we replace (2.2) - (2.4) by the system

$$\dot{T}_{i,j} + (1 - p_{i,j}^2) \dot{p}_{i,j} - a^2 \Delta_N T_{i,j} = 0 \quad (3.10)$$

and

$$\alpha \delta \dot{p}_{i,j} - \lambda^2 \delta^2 \Delta_N p_{i,j} = (\mu T_{i,j} + p_{i,j})(1 - p_{i,j}^2) \quad (3.11)$$

where $T_{i,j}$ and $p_{i,j}$ satisfy the discrete boundary conditions (3.5) and the symbol \cdot denotes differentiation with respect to t . Solutions of (3.10) and (3.11) obey the pointwise estimate that if $-1 \leq p_{i,j}(0^+) \leq 1$, $1 \leq i, j \leq N$, then $-1 \leq p_{i,j}(t) \leq 1$, $1 \leq i, j \leq N$ for all $t > 0$. This system also satisfies estimates which are the exact analogue of those obtained in section (2). Corresponding to (2.8) we have the identity

$$[T + p - p^3/3, 1](t) \equiv h_0 \quad (3.12)$$

while the analogues of (2.11) and (2.15) are the inequalities

$$\frac{d}{dt} \left(\left[\left(\frac{\mu T^2}{2\delta} + \frac{(1-p^2)^2}{4\delta} \right), 1 \right] - \frac{\lambda^2 \delta}{2} [p, \Delta_N p] \right) (t) = -\alpha [p_t, p_t](t) + \frac{\mu a^2}{\delta} [T, \Delta_N T](t), \quad (3.13)$$

and

$$\begin{aligned} & \frac{d}{dt} \left(\left[\frac{\mu T^2}{2\delta} + \frac{(1-p^2)^2}{4\delta}, 1 \right] - \frac{\lambda^2 \delta}{2} [p, \Delta_N p] - \frac{\alpha a^2}{2} [T, \Delta_N T] \right) (t) \\ & \leq -\frac{\alpha}{2} ([p_t, p_t] + [T_t, T_t]) (t) + \frac{\mu a^2}{\delta} [T, \Delta_N T] (t). \end{aligned} \quad (3.14)$$

We spare the reader the detailed analogues of (2.39), (2.41), and (2.45). Suffice it to say if we replace terms like $\int_{-1/2}^{1/2} \int_{-1/2}^{1/2} f(x, y, t) dx dy$ by $[f, 1](t)$ and terms like $\int_{-1/2}^{1/2} \int_{-1/2}^{1/2} |\nabla f|^2(x, y, t) dx dy$ by $-[f, \Delta_N f](t)$, then the resulting inequalities go through intact.

When doing simulations with the discrete system (3.10) and (3.11) we link the small parameter δ to the grid $dx = dy = 1/N$; specifically we set

$$\delta = 1/N \quad (3.15)$$

and restrict our attention to initial data satisfying

$$N \left[\frac{\mu T^2}{2} + \frac{(1-p^2)^2}{4}, 1 \right] (0^+) - \frac{\lambda^2}{2N} [p, \Delta_N p] (0^+) - \frac{\alpha a^2}{2} [T, \Delta_N T] (0^+) = 0(1). \quad (3.16)$$

Interesting data satisfying (3.16) may be achieved in a variety of ways. Below we outline one such construction. We let Γ be a simple closed curve of finite arclength $L(\Gamma)$ which is wholly contained in Ω , \mathcal{S}_c be all cells $\Omega_{i,j}$ which are cut by Γ , \mathcal{S}_I be all cells $\Omega_{i,j}$ which are interior to the region surrounded by Γ , and \mathcal{S}_E be those cells $\Omega_{i,j}$ which are in Ω and exterior to Γ . For definiteness we choose

$$p_{i,j}(0^+) = \begin{cases} 1, & \text{if } \Omega_{i,j} \in \mathcal{S}_E \\ p_{i,j}^* \in [-1, 1], & \text{if } \Omega_{i,j} \in \mathcal{S}_c \\ -1, & \text{if } \Omega_{i,j} \in \mathcal{S}_I \end{cases} \quad (3.17)$$

and

$$T_{i,j}(0^+) = h_0 - [(p - p^3/3)(0^+), 1] = 0(1/N^{1/2}). \quad (3.18)$$

Such data models the situation where initially water surrounds ice. With such data it is easily checked that $N[T^2(0^+), 1] = 0(1)$, while $N[(1 - p^2)^2, 1]$ and $-\frac{1}{N}[p, \Delta_N p]$ are $0(L(\Gamma))$ and thus the functional defined in (3.16) is $0(1)$ independently of N . For such data we note that $\delta^3[\Delta_N p, \Delta_N p] = \frac{1}{N^3}[\Delta_N p, \Delta_N p]$ is also $0(L(\Gamma))$ independently of N . The last functional enters in the discrete analogue of (2.68).

The simulations we present were all run with the following parameters:

$$(\alpha, \lambda, a, \mu) = (1, 1, 1, 1). \quad (3.19)$$

We chose $N = 200$

$$dx = dy = \delta = 1/200 \quad (3.20)$$

and

$$dt = dx/10. \quad (3.21)$$

With these parameters

$$m1 \stackrel{def}{=} dt/(dx)^2 = 10 \quad (3.22)$$

and

$$m2 \stackrel{def}{=} \frac{\delta dt}{(dx)^2} = \frac{1}{10} \quad (3.23)$$

and thus we were compelled to use an implicit integration scheme for (3.10). Our choice was ADI working with the fixed tri-diagonal matrix whose non-zero entries are

$$A(i, i) = \begin{cases} 1 + m1 & \text{if } i = 1 \text{ and } i = 200 \\ 1 + 2m1 & \text{if } 2 \leq i \leq 199 \end{cases} \quad (3.24)$$

and

$$A(i, i + 1) = A(i + 1, i) = -m1, \quad 1 \leq i \leq 199. \quad (3.25)$$

We used a first order explicit Euler scheme on (3.11).

Our first simulation was run with the following choice of initial data:

$$p_{i,j}(0^+) = \begin{cases} 1, & \text{if } \begin{cases} \{101 + i \leq j \leq 200 \text{ and } 1 \leq i \leq 99\} \text{ or} \\ \{302 - i \leq j \leq 200 \text{ and } 102 \leq i \leq 200\} \text{ or} \\ \{1 \leq j \leq 100 - i \text{ and } 1 \leq i \leq 99\} \text{ or} \\ \{1 \leq j \leq -101 + i \text{ and } 102 \leq i \leq 200\}, \end{cases} \\ 0, & \text{if } \begin{cases} \{100 + i = j \text{ and } 1 \leq i \leq 100\} \text{ or} \\ \{301 - i = j \text{ and } 101 \leq i \leq 200\} \text{ or} \\ \{101 - i = j \text{ and } 1 \leq i \leq 100\} \text{ or} \\ \{-100 + i = j \text{ and } 101 \leq i \leq 200\}, \end{cases} \\ -1, & \text{if } \begin{cases} \{102 - i \leq j \leq 99 + i \text{ and } 2 \leq i \leq 100\} \text{ or} \\ \{-99 + i \leq j \leq 300 - i \text{ and } 101 \leq i \leq 199\}. \end{cases} \end{cases} \quad (3.26)$$

This data satisfies $[p, 1] = [p^3, 1] = 0$. We chose $h_0 = .05$ and $T_{i,j}(0^+) \equiv .05$, $1 \leq i, j \leq 200$.

The initially superheated square block of ice converged to a circular block centered at $(0, 0)$ with a radius consistent with the $\delta = 0^+$ asymptotics. Specifically, if we grant that the ‘‘equilibrium’’ curve separating the ice from the water is a circle of radius R , then the ‘‘Gibbs-Thomson’’ relation (1.6), with our choice of parameters, implies that

$$\frac{.005}{2^{1/2}R} = -T_M \quad (3.27)$$

while (3.12) implies that

$$T_M = .05 - [p - p^3/3, 1]. \quad (3.28)$$

Noting that $p_{i,j} \simeq -1$ inside of the circle and $p_{i,j} \simeq +1$ outside yields

$$[p - p^3/3, 1] = \frac{2}{3} (1 - 2\pi R^2) \quad (3.29)$$

and (3.27)-(3.29) imply that R satisfies

$$\frac{4\pi}{3} R^2 + \frac{.005}{2^{1/2}R} = \frac{2}{3} - .05. \quad (3.30)$$

(3.30) has one positive solution close to zero and a second larger solution $R = .3808$ which is the approximate equilibrium radius we obtain in our simulation. This radius, when substituted into the Gibbs-Thomson relation (3.27) yields a melt temperature consistent with what we observe. The results of this simulation are shown in Figure 1.

Our final simulation starts with 16 regularly spaced circular blocks of ice surrounded by water. Specifically

$$p_{i,j}(0^+) = \begin{cases} -1, & \text{if } \sqrt{(i-40m)^2 + (j-40n)^2} \leq 15 \\ & \text{and } 1 \leq m, n \leq 4 \\ 1, & \text{otherwise.} \end{cases} \quad (3.31)$$

For this data

$$[(p - p^3/3)(0^+), 1] = .2885. \quad (3.32)$$

We again chose h_0 to be .05 and at $t = 0$ took the uniform temperature field

$$T_{i,j}(0^+) = h_0 - [(p - p^3/3)(0^+), 1] = -.2385. \quad (3.33)$$

The solutions using this initial data converged to the same equilibrium as before but the approach was far more interesting. The solutions at times $t = .05, .17, .24, .4, .6, .7, 2, 3.3,$ and 12.5 are shown in Figures 2-10.

We also ran both sets of data with the “intermediate” model and obtained virtually identical results.

Explanation of Graphics

In Figures 1-10 the upper left frame is a contour plot of p and p is 1 in a neighbourhood of the boundaries $x = \pm 1/2$ and $y = \pm 1/2$ and the upper right hand frame is a contour plot of T . In Figure 1 the lower left plot shows p, T and $p - p^3/3 + T$ along the lines $x = 0, y = 0,$ and the 45° diagonal through the origin. These plots overlay one another because of the circular symmetry of the final state. In Figure 1 the bottom right plot again displays the same information. In Figures 2-10 the bottom left plot shows $p, T,$ and $p - p^3/3 + T$ along the 45° diagonal through the origin and the bottom right plot shows the same quantities on the line $x = 0$.







Figure 1:







Figure 2:







Figure 3:







Figure 4:







Figure 5:







Figure 6:







Figure 7:







Figure 8:







Figure 9:







Figure 10:

References

1. S.L. Wang, R.F. Sekerka, A.A. Wheeler, B.T. Murray, S.R. Coriell, R.J. Braun, and G.B. McFadden, "Thermodynamically-Consistent Phase-Field Models, *Physica D* 69 (1993) 189-200.
2. G.B. McFadden, A.A. Wheeler, R.J. Brown, S.R. Coriell and R.F. Sekerka, "Phase-Field Models for Anisotropic Interfaces," *Phys. Rev. E* 48 (1993) 2016-2024.
3. S.L. Wang and R.F. Sekerka, "Algorithms for Phase Field Computations of the Dendritic Operating State at Large Supercoolings," *J. Comp. Physics* 127 (1996) 110-117.
4. A.S. Almgren and R.F. Almgren, "Phase Field Instabilities and Adaptive Mesh Refinement," in *Modern Methods for Modeling Microstructure in Materials*, TMS/SIAM 1996.
5. G. Caginalp, "Stefan and Hele-Shaw Type Models as Asymptotic Limits of the Phase Field Equation," *Phys. Rev. A* 39 (1989) 5887-5896.
6. G. Caginalp, "Phase Field Models and Sharp Interface Limits: Some Differences in Subtle Situations," *Rocky Mountain J. Math.* 21 (1996), 2, 603-616.