## 21-301 Combinatorics Homework 3 Due: Monday, September 27

- 1. Suppose that  $A_1, A_2, \ldots, A_n \subseteq A$  and  $|A_i| = k$  for  $i = 1, 2, \ldots, n$  and that q is a positive integer. Show that if  $nq \left(1 \frac{1}{q}\right)^k < 1$  then the elements of A can be q-colored so that each  $A_i$  contains an element of each color.
- Let G = (V, E) be a graph on n vertices, with minimum degree δ > 1. Show that G contains a dominating set of size at most n<sup>1+log(δ+1)</sup>/<sub>δ+1</sub>.
  (S is a dominating set if every v ∉ S has a neighbor in S.)
  (Hint: Choose S<sub>1</sub> ⊆ V by placing v into S<sub>1</sub> with probability p. Let S<sub>2</sub> denote the vertices in V \ S<sub>1</sub> that are not adjacent to a vertex in S<sub>1</sub>. Choose p carefully and use 1 − p ≤ e<sup>-p</sup>.)
- 3. Prove that there is an absolute constant c > 0 with the following property. Let A be an  $n \times n$  matrix with pairwise distinct real entries. Then there is a permutation of the rows of A so that no column in the permuted matrix contains an increasing subsequence of length at least  $c\sqrt{n}$ .

The following inequalities might be useful:

$$\binom{n}{k} \le \left(\frac{ne}{k}\right)^k$$
 and  $1 + x \le e^x$  and  $n! \ge \left(\frac{n}{e}\right)^n$ .