

21-301 Combinatorics
Homework 5
Due: Wednesday, October 13

1. Subsets $A_i, B_i \subseteq [n]$, $i = 1, 2, \dots, m$ satisfy (i) $A_i \cap B_i = \emptyset$ for all i and (ii) $A_i \cap B_j \neq \emptyset$ for all $i \neq j$. Show that

$$\sum_{i=1}^m \frac{1}{\binom{|A_i|+|B_i|}{|A_i|}} \leq 1.$$

(Hint: Let π be a random permutation of $[n]$ and for disjoint sets A, B define the event $\mathcal{E}(A, B)$ by

$$\mathcal{E}(A, B) = \{\pi : \max\{\pi(a) : a \in A\} < \min\{\pi(b) : b \in B\}\}.$$

Show that the events $\mathcal{E}_i = \mathcal{E}(A_i, B_i)$, $i = 1, 2, \dots, m$ are disjoint.)

2. Let x_1, x_2, \dots, x_n be real numbers such that $x_i \geq 1$ for $i = 1, 2, \dots, n$. Let J be any open interval of width 2. Show that of the 2^n sums $\sum_{i=1}^n \varepsilon_i x_i$, ($\varepsilon_i = \pm 1$), at most $\binom{n}{\lfloor n/2 \rfloor}$ lie in J .

(Hint: For $A \subseteq [n]$ let $x_A = \sum_{i \in A} x_i - \sum_{i \notin A} x_i$. Let $\mathcal{A} = \{A : x_A \in J\}$. Use Sperner's lemma.)

3. Suppose that we two-color the edges of K_6 Red and Blue. Show that there are at least two monochromatic triangles.